arXiv:0905.2516v1 [math.CO] 15 May 2009

An approach to the structure of imprimitive graphs which are not multicovers

Bin Jia

Center for Combinatorics
LPMC, Nankai University
Tianjin 300071
P. R. China
jiabinqq@gmail.com

Abstract. Denote by $\mathcal{G}$ the set of triples $(\Gamma, X, \mathcal{B})$, where Γ is a finite X-symmetric graph of valency $val(\Gamma) \geq 1$, $\mathcal{B}$ is a nontrivial X-invariant partition of $V(\Gamma)$ such that $\Gamma_{\mathcal{B}}$, the quotient graph of Γ with respect to $\mathcal{B}$, is nonempty and Γ is not a multicover of $\Gamma_{\mathcal{B}}$. In this article, for any given X-symmetric graph Σ, we aim to give a sufficient and necessary condition for the existence of $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, such that $\Gamma_{\mathcal{B}} \cong \Sigma$ and Γ is (X, s)-arc-transitive for some integer $s \geq 1$. And in this condition, a practicable method will be given to construct such an (X, s)-arc-transitive graph Γ. Moreover, for any $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, we will show that there exists a finite integer $m \geq 1$, such that after m times of such constructing process, we can obtain an X-symmetric graph $\Gamma_{\mathcal{B}_m}$ from $\Gamma_{\mathcal{B}}$, such that either $\Gamma \cong \Gamma_{\mathcal{B}_m}$ or Γ is a multicover of $\Gamma_{\mathcal{B}_m}$.

Keywords: symmetric graph; s-arc-transitive graph; imprimitive graph; quotient graph; double-star graph.

1 Introduction and Notation

In this paper, all graphs related are assumed to be finite, simple, nonempty and undirected.

Let Σ be a regular graph with vertex set $V(\Sigma)$, edge set $E(\Sigma)$ and valency $\mathbb{b} \geq 1$. Let $\mathbb{\sigma} := (\sigma_0, \sigma_1, \ldots, \sigma_{\mathbb{l}})$ be an $(\mathbb{l}+1)$-sequence of $V(\Sigma)$, where $\mathbb{l} \geq 0$ is an integer. Then $\mathbb{\sigma}$ is called an $\mathbb{l}$-arc of Σ if and only if $\{\sigma_i, \sigma_{i+1}\} \in E(\Sigma)$ for $i = 0, 1, \ldots, \mathbb{l}-1$, and $\sigma_{i-1} \neq \sigma_{i+1}$ for $i = 1, 2, \ldots, \mathbb{l}-1$. Denote by $Arc_{\mathbb{l}}(\Sigma)$ the set of $\mathbb{l}$-arcs of Σ. In the case of $\mathbb{l} = 1$, we use *arc* in place of 1-arc, and $Arc(\Sigma)$ in place of $Arc_1(\Sigma)$. For any $\mathbb{\sigma} = (\sigma_0, \sigma_1, \ldots, \sigma_{\mathbb{l}}) \in Arc_{\mathbb{l}}(\Sigma)$ and two integers i, j such that $0 \leq i \leq j \leq \mathbb{l}$, let $\mathbb{\sigma}(i, j) := (\sigma_i, \sigma_{i+1}, \ldots, \sigma_j)$, then $\mathbb{\sigma}(i, j) \in Arc_{j-i}(\Sigma)$.

For any $\sigma \in V(\Sigma)$, let $Arc_{\mathbb{l}}(\Sigma, \sigma) := \{\mathbb{\sigma} \in Arc_{\mathbb{l}}(\Sigma) |\ \mathbb{\sigma}(0,0) = (\sigma)\}$. When $\mathbb{l} = 1$, we use $\Sigma[\sigma]$ in place of $Arc_1(\Sigma, \sigma)$. For a nonempty subset $\mathbb{S}$ of $Arc_{\mathbb{l}}(\Sigma, \sigma)$ and an integer i such that $0 \leq i \leq \mathbb{l}$, let $\mathbb{S}(i) := \{\mathbb{\sigma}(0, i) |\ \mathbb{\sigma} \in \mathbb{S}\} \subseteq Arc_i(\Sigma, \sigma)$. And for $\mathbb{\alpha} \in \mathbb{S}(i)$, let $\mathbb{S}_{\mathbb{\alpha}} := \{\mathbb{\sigma} \in \mathbb{S} |\ \mathbb{\sigma}(0, i) = \mathbb{\alpha}\}$. In the case of $\mathbb{l} \geq 1$, for any $(\sigma_0, \tau_0) \in \mathbb{S}(1)$, let $\mathbb{S}_{\tau_0} := \{(\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in Arc_{\mathbb{l}-1}(\Sigma) | (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}\} \cup \{(\tau_0, \sigma_0, \sigma_1, \cdots, \sigma_{\mathbb{l}-2}) \in Arc_{\mathbb{l}-1}(\Sigma) | (\sigma_0, \sigma_1, \cdots, \sigma_{\mathbb{l}-2}) \in \mathbb{S}(\mathbb{l}-2)\}$. Then $\mathbb{S}_{\tau_0} = \{(\tau_0)\}$ if and only if $\mathbb{l} = 1$.

Obviously, $\{Arc_{\mathbb{l}}(\Sigma, \sigma) | \sigma \in V(\Sigma)\}$, where $\mathbb{l} \geq 1$, is a partition of $Arc_{\mathbb{l}}(\Sigma)$, and this partition is nontrivial when $\mathbb{b} \geq 2$. So, it is natural for us to study $Arc_{\mathbb{l}}(\Sigma)$ through the investigation of $Arc_{\mathbb{l}}(\Sigma, \sigma)$ for $\sigma \in V(\Sigma)$.

Let $\sigma \in V(\Sigma)$, $\mathbb{r}$ be an integer such that $1 \leq \mathbb{r} \leq \mathbb{b}$. For $\mathbb{S} \subseteq Arc_{\mathbb{l}}(\Sigma, \sigma)$, where $\mathbb{l} \geq 0$, $\mathbb{S}$ is called an $\left\llbracket {\mathbb{r} \atop \mathbb{l}} \right\rrbracket$-star of Σ with center σ if and only if one of the following three cases occurs:

(1). $\mathbb{l} = 0$, $\mathbb{S} = \{(\sigma)\}$ is called a 0-star.

(2). $\mathbb{l} = 1$, and $|\mathbb{S}| = \mathbb{r}$.

(3). $\mathbb{l} \geq 2$, $\mathbb{S}(\mathbb{l}-1)$ is an $\left\llbracket {\mathbb{r} \atop \mathbb{l}-1} \right\rrbracket$-star of Σ, and for any $\mathbb{a} \in \mathbb{S}(\mathbb{l}-1)$, $|\mathbb{S}_{\mathbb{a}}| = \mathbb{r}-1$.

Clearly, for any $\mathbb{S} \subseteq Arc_{\mathbb{l}}(\Sigma, \sigma)$ and $\tau \in V(\Sigma)$ such that $(\sigma, \tau) \in \mathbb{S}(1)$, where $\mathbb{l} \geq 1$, if $\mathbb{S}$ is an $\left\llbracket {\mathbb{r} \atop \mathbb{l}} \right\rrbracket$-star of Σ, then $\mathbb{S}_\tau$ is an $\left\llbracket {\mathbb{r} \atop \mathbb{l}-1} \right\rrbracket$-star of Σ. Hereafter, for any $\left\llbracket {\mathbb{r} \atop \mathbb{l}} \right\rrbracket$-star $\mathbb{S}$ of Σ, we use $\mathfrak{C}_\Sigma(\mathbb{S})$ to denote the center of $\mathbb{S}$ in Σ.

Denote by $St^{\mathbb{r}}_{\mathbb{l}}(\Sigma)$ the set of $\left\llbracket {\mathbb{r} \atop \mathbb{l}} \right\rrbracket$-stars of Σ. In the case of $\mathbb{l} = 1$, we use $\mathbb{r}$-star and $St^{\mathbb{r}}(\Sigma)$ in place of $\left\llbracket {\mathbb{r} \atop 1} \right\rrbracket$-star and $St^{\mathbb{r}}_1(\Sigma)$, respectively.

Let Σ be an X-symmetric graph with valency $\mathbb{b} \geq 1$. For $\mathbb{S} \subseteq Arc_{\mathbb{l}}(\Sigma, \sigma)$, where $\mathbb{l} \geq 0$ is an integer, denote by $X_{\mathbb{S}}$ the set-wise stabilizer of X on $\mathbb{S}$, then $X_\sigma = X_{\mathbb{S}(0)} \geq X_{\mathbb{S}(1)} \geq \cdots \geq X_{\mathbb{S}(\mathbb{l}-1)} \geq X_{\mathbb{S}(\mathbb{l})} = X_{\mathbb{S}}$. For any integer s with $1 \leq s \leq \mathbb{l}$, $\mathbb{S}$ is said to be $(X_{\mathbb{S}}, s)$-arc-transitive if and only if $X_{\mathbb{S}}$ acts transitively on $\mathbb{S}(s)$.

An X-orbit S of $St^{\mathbb{r}}_{\mathbb{l}}(\Sigma)$ is said to be (X, s)-arc-transitive, where $1 \leq s \leq \mathbb{l}$, if and only if any $\mathbb{S} \in S$ is $(X_{\mathbb{S}}, s)$-arc-transitive. Clearly, if $\mathbb{S}$ is $(X_{\mathbb{S}}, s)$-arc-transitive for some integer $s \geq 2$, then $\mathbb{S}$ is $(X_{\mathbb{S}}, s-1)$-arc-transitive.

For $\mathbb{S}, \mathbb{T} \in St^{\mathbb{r}}_{\mathbb{l}}(\Sigma)$, where $\mathbb{l} \geq 1$, let $\mathfrak{C}_\Sigma(\mathbb{S}) = \sigma$, $\mathfrak{C}_\Sigma(\mathbb{T}) = \tau$. $(\mathbb{S}, \mathbb{T})$ is called a $\left\llbracket {\mathbb{r} \atop \mathbb{l}} \right\rrbracket$-double-star of Σ, if and only if $(\sigma, \tau) \in \mathbb{S}(1)$, $(\tau, \sigma) \in \mathbb{R}(1)$, $\mathbb{S}_\tau = \mathbb{T}(\mathbb{l}-1)$ and $\mathbb{T}_\sigma = \mathbb{S}(\mathbb{l}-1)$. We depict a $\left\llbracket {3 \atop 2} \right\rrbracket$-star and a $\left\llbracket {3 \atop 2} \right\rrbracket$-double-star in Figure 1.


a star

a double-star

Figure 1. A star and a double-star

Denote by $DSt^{\mathbb{r}}_{\mathbb{l}}(\Sigma)$ the set of $\llbracket {\mathbb{r} \atop \mathbb{l}} \rrbracket$-double-stars of Σ, where $\mathbb{l} \geq 1$. In the case of $\mathbb{l} = 1$, we use $\mathbb{r}$-double-star and $DSt^{\mathbb{r}}(\Sigma)$ in place of $\llbracket {\mathbb{r} \atop 1} \rrbracket$-double-star and $DSt^{\mathbb{r}}_{1}(\Sigma)$, respectively. A subset Θ of $DSt^{\mathbb{r}}_{\mathbb{l}}$ is said to be self-paired if and only if $(\mathbb{S}, \mathbb{T}) \in \Theta$ always implies that $(\mathbb{T}, \mathbb{S}) \in \Theta$.

For $1 \leq s \leq \mathbb{l}$, an X-orbit Θ of $DSt^{\mathbb{r}}_{\mathbb{l}}(\Sigma)$ is said to be (X, s)-arc-transitive, if and only if any $\mathbb{S} \in St(\Theta) := \{\mathbb{L}, \mathbb{R} | (\mathbb{L}, \mathbb{R}) \in \Theta\}$ is $(X_{\mathbb{S}}, s)$-arc-transitive. A self-paired $(X, 1)$-arc-transitive orbit Θ of $DSt^{\mathbb{r}}_{\mathbb{l}}(\Sigma)$ is said to be X-symmetric.

Let Γ be a finite X-symmetric graph with vertex set $V(\Gamma)$ and edge set $E(\Gamma)$. A partition $\mathcal{B}$ on $V(\Gamma)$ with $|\mathcal{B}| \geq 2$ is said to be X-invariant if and only if for any $B \in \mathcal{B}$ and $x \in X$, $B^x := \{\sigma^x | \sigma \in B\} \in \mathcal{B}$. In this case, the quotient graph $\Gamma_{\mathcal{B}}$ of Γ with respect to $\mathcal{B}$ is defined to be with vertex set $\mathcal{B}$, and for $B, C \in \mathcal{B}$, B adjacent to C if and only if there exists some $\sigma \in B$ adjacent to some $\tau \in C$. We always assume that $\Gamma_{\mathcal{B}}$ is nonempty, which implies that $\Gamma_{\mathcal{B}}$ is X-symmetric, and by [1], any $B \in \mathcal{B}$ is an independent set of Γ. For any $\sigma \in V(\Gamma)$, $B \in \mathcal{B}$, let $\Gamma(\sigma)$ be the neighborhood of σ in Γ, $\Gamma_{\mathcal{B}}(B)$ be the neighborhood of B in $\Gamma_{\mathcal{B}}$, $\Gamma(B) := \bigcup_{\iota \in B} \Gamma(\iota)$, $\Gamma_{\mathcal{B}}(\sigma) := \{C \in \mathcal{B} | \sigma \in \Gamma(C)\}$. For any $(B, C) \in Arc(\Gamma_{\mathcal{B}})$ and $\sigma \in V(\Gamma)$, denote by $\Gamma[B, C]$ the bipartite subgraph of Γ induced by $(\Gamma(B) \cap C) \cup (B \cap \Gamma(C))$. Let $v := |B|$, $k := |\Gamma(B) \cap C|$, $r := |\Gamma_{\mathcal{B}}(\sigma)|$, $b := val(\Gamma_{\mathcal{B}})$, $d := val(\Gamma[B, C])$, then $p(\Gamma, X, \mathcal{B}) := (v, k, r, b, d)$ is independent of the choice of $\sigma \in V(\Gamma)$ and $(B, C) \in Arc(\Gamma_{\mathcal{B}})$. When $v = 1$, we say $\mathcal{B} = 1$ is trivial. Otherwise, $v \geq 2$ and we say $\mathcal{B} > 1$ is nontrivial. When $\mathcal{B} > 1$ and $v = k \geq 2$, Γ is called a multicover of $\Gamma_{\mathcal{B}}$. A multicover Γ of $\Gamma_{\mathcal{B}}$ is called a cover of $\Gamma_{\mathcal{B}}$ if and only if $d = 1$.

In this paper, we mainly discuss the case of $v > k \geq 1$, that is, $\mathcal{B} > 1$ and Γ is not a multicover of $\Gamma_{\mathcal{B}}$. The following is a summary of our main results:

Theorem 1.1 *Let Σ be a finite X-symmetric graph with valency $\mathbb{b} \geq 2$, and let $s \geq 1$ be an integer. Then there exists a finite (X, s)-arc-transitive graph Γ with a nontrivial X-invariant partition $\mathcal{B}$ on $V(\Gamma)$, such that $\Gamma_{\mathcal{B}} \cong \Sigma$ and $1 \leq r \leq \mathbb{b} - 1$, if and only if one of the following cases occurs:*

(a). *$s = 1$, there exists an X-symmetric orbit Θ on $DSt^{r}(\Sigma)$.*

(b). *$s \geq 2$, $r = 1$, and for any $\{\sigma, \tau\} \in E(\Sigma)$, there exists a bipartite $(X_{\{\sigma,\tau\}}, s)$-arc-transitive graph Ψ such that for any $\{\psi, \delta\} \in E(\Psi)$, $X_{\psi} < X_{(\sigma,\tau)}$, $X_{\{\psi,\delta\}} < X_{\{\sigma,\tau\}}$ and $X_{\{\psi,\delta\}} \setminus X_{(\sigma,\tau)} \neq \emptyset$.*

(c). *$s \geq 2$, $\mathbb{b} \geq 3$, $r \geq 2$, there exists a self-paired (X, s)-arc-transitive orbit Θ on $DSt^{r}_{\mathbb{l}}(\Sigma)$, where $\mathbb{l} \geq s$ is a finite integer, such that for any $(\mathbb{S}, \mathbb{T}) \in \Theta$ with $\mathfrak{C}_{\Sigma}(\mathbb{S}) = \sigma$ and $\mathfrak{C}_{\Sigma}(\mathbb{T}) = \tau$, $X_{\mathbb{S}} \cap X_{\tau} = X_{\mathbb{T}} \cap X_{\sigma}$.*

More details about the case of $s = 1$ will been seen in Section 2, where we will discuss the double-stars and the double-star graph construction. As for the case of $s \geq 2$, we will discuss the $\llbracket {\mathbb{r} \atop \mathbb{l}} \rrbracket$-double-stars and the corresponding graph construction in Section 3.

2 Double-star graphs

In this section, we aim to find a practicable method to approach Γ for a given quotient $\Gamma_{\mathcal{B}}$. The method is mainly based on the following double-star graph construction.

Construction 2.1 *Let* Σ *be a finite* X*-symmetric graph with valency* $\mathbb{b} \geq 1$*. If* Σ *contains an* X*-symmetric orbit* Θ *on* $DSt^{\mathbb{r}}(\Sigma)$*, where* $1 \leq \mathbb{r} \leq \mathbb{b}$*, then the double-star graph* $\Pi(\Sigma, \Theta)$ *of* Σ *with respect to* Θ *is defined to be the graph with vertex set* $St(\Theta)$*, and edge set* $\{\{\mathbb{S}, \mathbb{T}\} | (\mathbb{S}, \mathbb{T}) \in \Theta\}$*.*

For the sake of convenient, we denote by $\mathbb{G}$ the set of the triples (Σ, X, Θ), where Σ is a finite X-symmetric graph with valency $\mathbb{b} \geq 2$, Θ is an X-symmetric orbit on $DSt^{\mathbb{r}}_{\mathbb{l}}(\Sigma)$, where $\mathbb{l} \geq 1$ and $1 \leq \mathbb{r} \leq \mathbb{b} - 1$. And for any $(\Sigma, X, \Theta) \in \mathbb{G}$, let $p(\Sigma, X, \Theta) = (\mathbb{b}, \mathbb{r}, \mathbb{l})$. In the case of $\mathbb{l} = 1$, we use $(\mathbb{b}, \mathbb{r})$ short for $(\mathbb{b}, \mathbb{r}, 1)$.

Also, we use $\mathcal{G}$ to denote the set of triples $(\Gamma, X, \mathcal{B})$, where Γ is a finite X-symmetric graph, $\mathcal{B}$ is a nontrivial X-invariant partition of $V(\Gamma)$ such that $\Gamma_{\mathcal{B}}$ is nonempty and Γ is not a multicover of $\Gamma_{\mathcal{B}}$.

Let Γ be an X-symmetric graph with an X-invariant partition $\mathcal{B}$ on $V(\Gamma)$, such that $val(\Gamma_{\mathcal{B}}) \geq 1$. Then for any $\sigma \in V(\Gamma)$, $\Gamma_{\mathcal{B}}[\sigma] := \{(B, C) \in Arc(\Gamma_{\mathcal{B}}) | \sigma \in B\}$ is an r-star of $\Gamma_{\mathcal{B}}$ with center $B \in \mathcal{B}$ such that $\sigma \in B$. Moreover, for any $\mathbb{r}$-star $\mathbb{S}$ of $\Gamma_{\mathcal{B}}$ with center $B \in \mathcal{B}$, let $\mathfrak{C}_{\Gamma}(\mathbb{S}) := \bigcap_{(B,C) \in \mathbb{S}} (B \cap \Gamma(C))$. Then, for any $\sigma \in B \in \mathcal{B}$, it is clear that $\sigma \in \mathfrak{C}_{\Gamma}(\Gamma_{\mathcal{B}}[\sigma]) \subseteq B$, and that $\mathfrak{C}_{\Gamma}(\Gamma_{\mathcal{B}}[\sigma]) = B$ if and only if $r = b$ (and $v = k$), that is, $\mathcal{B} = 1$ or Γ is a multicover of $\Gamma_{\mathcal{B}}$.

Theorem 2.2 *Let* $(\Sigma, X, \Theta) \in \mathbb{G}$*,* $\Pi = \Pi(\Sigma, \Theta)$*,* $S = St(\Theta)$*,* $\Pi = \Pi(\Sigma, \Theta)$*,* $\mathcal{S} = \{S_\sigma | \sigma \in V(\Sigma)\}$*, where* $S_\sigma := \{\mathbb{S} \in S | \mathfrak{C}_{\Sigma}(\mathbb{S}) = \sigma\}$ *for any* $\sigma \in V(\Sigma)$*. Then* $(\Pi, X, \mathcal{S}) \in \mathcal{G}$ *such that* $\Pi_{\mathcal{S}} \cong \Sigma$*. Moreover, for any* $\mathbb{S} \in S = V(\Pi)$*,* $\Pi_{\mathcal{S}}[\mathbb{S}] = \{(S_\sigma, S_\tau) | (\sigma, \tau) \in \mathbb{S}\}$ *and* $\mathfrak{C}_{\Pi}(\Pi_{\mathcal{S}}[\mathbb{S}]) = \{\mathbb{S}\}$*.*

Proof. It is not difficult to check that $\mathcal{S}$ is a nontrivial X-invariant partition of $V(\Pi)$. Note that $\sigma \leftrightarrow S_\sigma$, where $\sigma \in V(\Sigma)$, is a bijection between Σ and $\Pi_{\mathcal{S}}$, so $\Pi_{\mathcal{S}} \cong \Sigma$.

Let $p(\Sigma, X, \Theta) = (\mathbb{b}, \mathbb{r}, \mathbb{l})$ and let $p(\Pi, X, \mathcal{S}) = (v, k, r, b, d)$, then $b = \mathbb{b} \leq \mathbb{r} - 1 = r - 1$ and hence $(\Pi, X, \mathcal{S}) \in \mathcal{G}$.

For any $\mathbb{S} \in S = V(\Pi)$, we have $\Pi_{\mathcal{S}}[\mathbb{S}] = \{(S_\sigma, S_\tau) |\ \mathbb{S} \in S_\sigma \in \mathcal{S}, S_\tau \in \Pi_{\mathcal{S}}(\mathbb{S})\} = \{(S_\sigma, S_\tau) | \mathbb{S} \in S_\sigma \in \mathcal{S}$ and $\exists \mathbb{R} \in S_\tau$ such that $(\mathbb{S}, \mathbb{R}) \in \Theta\} = \{(S_\sigma, S_\tau) | (\sigma, \tau) \in \mathbb{S}\}$.

For any $\mathbb{S} \in S_\sigma \in \mathcal{S}$, where $\sigma \in V(\Sigma)$, it is clear that $\mathbb{S} \in \mathfrak{C}_{\Pi}(\mathbb{S}) \subset S_\sigma$. For any $\mathbb{L} \in \mathfrak{C}_{\Pi}(\Pi_{\mathcal{S}}[\mathbb{S}])$, we have $\Pi_{\mathcal{S}}[\mathbb{S}] \subseteq \Pi_{\mathcal{S}}[\mathbb{L}]$. For any $(\sigma, \tau) \in \mathbb{S}$, then $(S_\sigma, S_\tau) \in \Pi_{\mathcal{S}}[\mathbb{S}] \subseteq \Pi_{\mathcal{S}}[\mathbb{L}] \Rightarrow \mathbb{L} \in S_\sigma \in \mathcal{S}$ and there exists $\mathbb{R} \in S_\tau$ such that $(\mathbb{L}, \mathbb{R}) \in \Theta \Rightarrow (\sigma, \tau) \in \mathbb{L}$, it follows that $\mathbb{S} \subseteq \mathbb{L}$. Note that $|\mathbb{S}| = |\mathbb{L}| = \mathbb{r}$, hence $\mathbb{L} = \mathbb{R}$, and so we have $\mathfrak{C}_{\Pi}(\Pi_{\mathcal{S}}[\mathbb{S}]) = \{\mathbb{S}\}$. ∎

Let $\mathcal{B}_1$ and $\mathcal{B}_2$ be two X-invariant partitions on $V(\Gamma)$. Denote by $\mathcal{B}_2 \leq \mathcal{B}_1$ (or $\mathcal{B}_1 \geq \mathcal{B}_2$) that $\mathcal{B}_2$ is a refinement of $\mathcal{B}_1$, and by $\mathcal{B}_2 < \mathcal{B}_1$ (or $\mathcal{B}_1 > \mathcal{B}_2$) that $\mathcal{B}_2$ is a proper refinement of $\mathcal{B}_1$. For two X-invariant partitions $\mathcal{B}_1$, $\mathcal{B}_2$ of $V(\Gamma)$ such that $\mathcal{B}_1 \geq \mathcal{B}_2$, let $\mathcal{B}_1/\mathcal{B}_2 := \{\{B_2 \in \mathcal{B}_2 | B_2 \subseteq B_1\} | B_1 \in \mathcal{B}_1\}$, then it is clear that $\mathcal{B}_1/\mathcal{B}_2$ is an X-invariant partition on $V(\Gamma_{\mathcal{B}_2})$, such that $(\Gamma_{\mathcal{B}_2})_{\mathcal{B}_1/\mathcal{B}_2} \cong \Gamma_{\mathcal{B}_1}$, and that $\mathcal{B}_1/\mathcal{B}_2$ is nontrivial if and only if $\mathcal{B}_1 > \mathcal{B}_2$.

The following lemma shows that there exists a natural refinement of $\mathcal{B}$ for any $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$. This can be seen as an access to the structure of Γ.

Lemma 2.3 *Let* $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ *with* $p(\Gamma, X, \mathcal{B}) = (v, k, r, b, d)$, $S_{(0,1)} := \{\Gamma_{\mathcal{B}}[\tau] | \tau \in V(\Gamma)\}$, $\Theta_{(0,1)} := \{(\Gamma_{\mathcal{B}}[\tau], \Gamma_{\mathcal{B}}[\sigma]) | (\tau, \sigma) \in Arc(\Gamma)\}$, *then* $S_{(0,1)}$ *is an* X*-symmetric orbit on* $St^r(\Gamma_{\mathcal{B}})$ *while* $\Theta_{(0,1)}$ *is an* X*-symmetric orbit on* $DSt^r(\Gamma_{\mathcal{B}})$ *such that* $S_{(0,1)} = St(\Theta_{(0,1)})$. *Let* $\mathcal{B}_1 = \mathfrak{C}_\Gamma(S_{(0,1)}) := \{\mathfrak{C}_\Gamma(\mathbb{S}) | \mathbb{S} \in S_{(0,1)}\}$. *Then all of the following cases hold:*

- **(1).** $\mathcal{B}_1$ *is an* X*-invariant partition of* $V(\Gamma)$, *such that* $\mathcal{B} > \mathcal{B}_1 \geq 1$. *Let* $p(\Gamma, X, \mathcal{B}_1) = (v_1, k_1, r_1, b_1, d_1)$, *then* $v_1 = |\mathfrak{C}_\Gamma(\mathbb{S})| \geq 1$ *for any* $\mathbb{S} \in S_{(0,1)}$. *Further,* $v_1|(v, k)$, $r|(r_1, b_1)$, $d_1|d$, $v_1 r_1 = b_1 k_1$, $r_1 d_1 = rd = val(\Gamma)$.
- **(2).** $(\Gamma_{\mathcal{B}_1}, X, \mathcal{B}/\mathcal{B}_1) \in \mathcal{G}$ *with* $p(\Gamma_{\mathcal{B}_1}, X, \mathcal{B}/\mathcal{B}_1) = (v/v_1, k/v_1, r, b, b_1/r)$, *such that* $(\Gamma_{\mathcal{B}_1})_{\mathcal{B}/\mathcal{B}_1} \cong \Gamma_{\mathcal{B}}$ *and* $\Gamma_{\mathcal{B}_1} \cong \Pi(\Gamma_{\mathcal{B}}, \Theta_{(0,1)})$.

Moreover, one of the following cases occurs:

- **(a).** $v_1 = k_1 = 1$, $\Gamma \cong \Pi(\Gamma_{\mathcal{B}}, \Theta_{(0,1)})$, *and* X *acts faithfully on* $\mathcal{B}$ *if and only if* X *acts faithfully on* $V(\Gamma)$.
- **(b).** $v_1 = k_1 \geq 2$, *and* Γ *is a multicover of* $\Gamma_{\mathcal{B}_1}$.
- **(c).** $v_1 > k_1 \geq 1$, *and* $(\Gamma, X, \mathcal{B}_1) \in \mathcal{G}$.

Proof. It is not difficult to check that $S_{(0,1)}$ is an X-symmetric orbit on $St^r(\Gamma_{\mathcal{B}})$, $\Theta_{(0,1)}$ is a self-paired X-symmetric orbit on $DSt^r(\Gamma_{\mathcal{B}})$, $S_{(0,1)} = St(\Theta_{(0,1)})$.

- **(1).** For any $(B, C) \in \mathbb{S} = \Gamma_{\mathcal{B}}[\sigma] \in S_{(0,1)}$, where $\sigma \in B \in \mathcal{B}$, by the previous analysis, $\emptyset \subset \{\sigma\} \subseteq \mathfrak{C}_\Gamma(\mathbb{S})$. Additionally, $\mathfrak{C}_\Gamma(\mathbb{S}) \subseteq \Gamma(C) \cap B$, hence $1 \leq |\mathfrak{C}_\Gamma(\mathbb{S})| \leq k$. For any $x \in X$, since $\mathfrak{C}_\Gamma(\mathbb{S})^x = \mathfrak{C}_\Gamma(\mathbb{S}^x)$, hence $\mathcal{B}_1$ is an X-invariant partition of $V(\Gamma)$ and is a proper refinement of $\mathcal{B}$, such that $1 \leq \mathcal{B}_1 < \mathcal{B}$. Thus $v_1 = |\mathfrak{C}_\Gamma(\mathbb{S})| \in [1, k]$ is independent of the choice of $\mathbb{S} \in S_{(0,1)}$ and $v_1|v$.

  Obviously, $X_{\Gamma(C)\cap B}$ acts transitively on $\Gamma(C) \cap B$. Consider $\{\mathfrak{C}_\Gamma(\Gamma_{\mathcal{B}}[\sigma])| \sigma \in \Gamma(C) \cap B\}$, easy checking shows that it is an $X_{\Gamma(C)\cap B}$-invariant partition of $\Gamma(C) \cap B$, so $v_1|k$, and hence $v_1|(v, k)$.

It is clear that $v_1 r_1 = b_1 k_1$ and $r_1 d_1 = rd = val(\Gamma)$. For any $\sigma \in B$, as Γ is X-symmetric, $X_\sigma \cap X_{\Gamma(\sigma)\cap C}$ acts transitively on $\Gamma(\sigma)\cap C$. Moreover, it is easy to check that $\{\Gamma(\sigma)\cap C_1 | C_1 \in \mathcal{B}_1, C_1 \subseteq C\}$ is an $X_\sigma \cap X_{\Gamma(\sigma)\cap C}$-invariant partition of $\Gamma(\sigma)\cap C$, hence $r_1/r = d/d_1 = |\{\Gamma(\sigma)\cap C_1 | C_1 \in \mathcal{B}_1, C_1 \subseteq C\}| \in \mathbb{Z}^+$. Further, for $B_1 \in \mathcal{B}_1$ such that $B_1 \subseteq B \in \mathcal{B}$, it is not difficult to see that $\{\{C_1 \in \Gamma_{\mathcal{B}_1}(B_1) | C_1 \subseteq C\} | C \cap \Gamma(B_1) \neq \emptyset\}$ is an X_{B_1}-invariant partition of an X_{B_1}-transitive set $\Gamma_{\mathcal{B}_1}(B_1)$. Thus $b_1 = |\Gamma_{\mathcal{B}_1}(B_1)|$ is divided by $r = |\{\{C_1 \in \Gamma_{\mathcal{B}_1}(B_1) | C_1 \subseteq C\} | C \cap \Gamma(B_1) \neq \emptyset\}|$, and hence $r | (r_1, b_1)$.

(2). Easy checking shows that $(\Gamma_{\mathcal{B}_1}, X, \mathcal{B}/\mathcal{B}_1) \in \mathcal{G}$, that $(\Gamma_{\mathcal{B}_1})_{\mathcal{B}/\mathcal{B}_1} \cong \Gamma_{\mathcal{B}}$ and that $\Gamma_{\mathcal{B}_1} \cong \Pi(\Gamma_{\mathcal{B}}, \Theta_{(0,1)})$. Denote by $(v_{01}, k_{01}, r_{01}, b_{01}, d_{01})$ the parameters of $(\Gamma_{\mathcal{B}_1}, X, \mathcal{B}/\mathcal{B}_1)$, then for any $B \in \mathcal{B}$, as $\{B_1 \in \mathcal{B}_1 | B_1 \subseteq B\}$ is an X_B-invariant partition of B and X_B acts transitively on B, we have $v_{01} = |\{B_1 \in \mathcal{B}_1 | B_1 \subseteq B\}| = v/v_1$. For $(B, C) \in Arc(\Gamma_{\mathcal{B}})$, since $X_{(B,C)}$ acts transitively on $B \cap \Gamma(C)$, and $\{B_1 \in \mathcal{B} | B_1 \subseteq B \cap \Gamma(C)\}$ is an $X_{(B,C)}$-invariant partition of $B \cap \Gamma(C)$, so $k_{01} = |\{B_1 \in \mathcal{B} | B_1 \subseteq B \cap \Gamma(C)\}| = k/v_1$. It is clear that $r_{01} = r$, $b_{01} = b = val(\Gamma_{\mathcal{B}})$. For $B_1 \in \mathcal{B}_1$, by the analysis in **(1)**, $\{\{C_1 \in \Gamma_{\mathcal{B}_1}(B_1) | C_1 \subseteq C\} | C \cap \Gamma(B_1) \neq \emptyset\}$ is an X_{B_1}-invariant partition of an X_{B_1}-transitive set $\Gamma_{\mathcal{B}_1}(B_1)$, so $b_1 = |\Gamma_{\mathcal{B}_1}(B_1)| = |\{\{C_1 \in \Gamma_{\mathcal{B}_1}(B_1) | C_1 \subseteq C\} | C \cap \Gamma(B_1) \neq \emptyset\}| \cdot |\{D_1 \in \Gamma_{\mathcal{B}_1}(B_1) | D_1 \subseteq D\}| = r \cdot d_{01}$, where $D \in \mathcal{B}$ such that $D \cap \Gamma(B_1) \neq \emptyset$. Hence we have $d_{01} = b_1/r$.

As $1 \leq k_1 \leq v_1$, one of the following cases occurs:

(a). $v_1 = k_1 = 1$. In this case $\mathcal{B}_1 = 1$ and hence $\Gamma \cong \Gamma_{\mathcal{B}_1} \cong \Pi(\Gamma_{\mathcal{B}}, \Theta)$. On the one hand, it is clear that $\bigcap_{\gamma \in V(\Gamma)} X_\gamma \leq \bigcap_{B \in \mathcal{B}} X_B$. On the other hand, $\bigcap_{B \in \mathcal{B}} X_B = \bigcap_{B \in \mathcal{B}} (\bigcap_{C \in \Gamma_{\mathcal{B}}(B)} X_{(B,C)}) \leq \bigcap_{B \in \mathcal{B}} (\bigcap_{\gamma \in B} X_{\Gamma[\gamma]}) = \bigcap_{\gamma \in V(\Gamma)} X_{\mathfrak{C}_\Gamma(\Gamma[\gamma])}$. By Theorem 2.2, $\mathfrak{C}_\Gamma(\Gamma[\gamma]) = \{\gamma\}$, so $\bigcap_{B \in \mathcal{B}} X_B \leq \bigcap_{\gamma \in V(\Gamma)} X_\gamma$. It follows that $\bigcap_{B \in \mathcal{B}} X_B = \bigcap_{\gamma \in V(\Gamma)} X_\gamma$, and hence X acts faithfully on $\mathcal{B}$ if and only if X acts faithfully on $V(\Gamma)$.

(b). $v_1 = k_1 \geq 2$. In this case $\mathcal{B}_1$ is nontrivial and $v_1 = k_1$, hence Γ is a multicover of $\Gamma_{\mathcal{B}_1}$.

(c). $v_1 > k_1 \geq 1$. In this case $\mathcal{B}_1$ is nontrivial and $k_1 < v_1$, hence $(\Gamma, X, \mathcal{B}_1) \in \mathcal{G}$.

∎

In the following theorem, we will show that for any $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, we can construct the minimum imprimitive graph $\Gamma_{\mathcal{B}_m}$ from $\Gamma_{\mathcal{B}}$ through double-star graph construction, such that $(\Gamma, X, \mathcal{B}_m) \in \mathcal{G}$ and either $\Gamma \cong \Gamma_{\mathcal{B}_m}$ or Γ is a multicover of $\Gamma_{\mathcal{B}_m}$. Meanwhile, we will characterize the structure of Γ by counting some important parameters.

Theorem 2.4 *Let* $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ *with* $p(\Gamma, \mathcal{B}) = (v, k, r, b, d)$, *and let*

$$(v, k) = p_1^{n_1} \cdots p_t^{n_t},\ n_i \in \mathbb{Z}^+\ (i = 1, \cdots, t),$$

where $p_1 < \cdots < p_t$ *are primes. Then there exists an* $(m+1)$*-sequence, where* $1 \le m \le \sum_{i=1}^{t} n_i + 1$, *of* X*-invariant partitions*

$$\mathcal{B} = \mathcal{B}_0 > \mathcal{B}_1 > \cdots > \mathcal{B}_{m-1} > \mathcal{B}_m \ge 1$$

on $V(\Gamma)$ *with* $p(\Gamma, \mathcal{B}_i) = (v_i, k_i, r_i, b_i, d_i)$ *for* $i = 0, \cdots, m$, *such that all the following statements hold:*

(1). $(\Gamma, X, \mathcal{B}_i) \in \mathcal{G}$, $v_i \ge 2$ *and* $1 \le k_i \le v_i - 1$ *for* $i = 0, 1, \ldots, m-1$.

(2). $v_{i+1} | (v_i, k_i)$, $r_i | (r_{i+1}, b_{i+1})$ *and* $d_{i+1} | d_i$ *for* $i = 0, 1, \ldots, m$. *Moreover,* $v_i r_i = b_i k_i$, $r_i d_i = val(\Gamma)$ *for* $i = 0, 1, \ldots, m$.

(3). *Let* $S_{(i,i+1)} := \{\Gamma_{\mathcal{B}_i}[\tau] | \tau \in V(\Gamma)\}$, $\Theta_{(i,i+1)} := \{(\Gamma_{\mathcal{B}_i}[\tau], \Gamma_{\mathcal{B}_i}[\sigma]) | (\tau, \sigma) \in Arc(\Gamma)\}$, *then* $S_{(i,i+1)} = St(\Theta_{(i,i+1)})$ *is an* X*-symmetric orbit on* $St^{r_i}(\Gamma_{\mathcal{B}_i})$ *and* $\Theta_{(i,i+1)}$ *is a self-paired* X*-symmetric orbit on* $DSt^{r_i}(\Gamma_{\mathcal{B}_i})$ *for* $i = 0, 1, \ldots, m-1$. *Moreover,* $\mathcal{B}_{i+1} = \mathfrak{C}_\Gamma(S_{(i,i+1)})$ *and* $\Gamma_{\mathcal{B}_{i+1}} \cong \Pi(\Gamma_{\mathcal{B}_i}, \Theta_{(i,i+1)})$ *for* $i = 0, 1, \ldots, m-1$.

(4). $(\Gamma_{\mathcal{B}_j}, X, \mathcal{B}_i/\mathcal{B}_j) \in \mathcal{G}$ *with parameters* $(v_i/v_j, k_i/v_j, r_i, b_i, b_j/r_i)$ *such that* $(\Gamma_{\mathcal{B}_j})_{\mathcal{B}_i/\mathcal{B}_j} \cong \Gamma_{\mathcal{B}_i}$, *for* $0 \le i < j \le m$.

(5). *Either of the following two cases occurs:*

- **(5.1).** $v_m = k_m = 1$, $\mathcal{B}_m = 1$ *and* $\Gamma \cong \Pi(\Gamma_{\mathcal{B}_{m-1}}, \Theta_{(m-1,m)})$. *Additionally,* X *is faithful on* $\mathcal{B}_i$ *if and only if* X *if faithful on* $V(\Gamma)$, *for* $0 \le i \le m$.
- **(5.2).** $v_m = k_m \ge 2$, $\mathcal{B}_m$ *is a nontrivial* X*-invariant partition of* $V(\Gamma)$, *such that* Γ *is a multicover of* $\Gamma_{\mathcal{B}_m}$. *Additionally,* X *is faithful on* $\mathcal{B}_i$ *if and only if* X *is faithful on* $\mathcal{B}_j$, *for* $0 \le i, j \le m$.

Proof. For any integer $m \ge 1$, denote by $\mathcal{G}(m)$ the set of triples $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ with an $(m+1)$-sequence of X-invariant partitions

$$\mathcal{B} = \mathcal{B}_0 > \cdots > \mathcal{B}_m \ge 1$$

on $V(\Gamma)$ such that cases **(1)**-**(4)** hold. Then by Lemma 2.3, $\mathcal{G}(1) = \mathcal{G}$, that is, for any given $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, there always exists some $m \ge 1$ such that $(\Gamma, X, \mathcal{B}) \in \mathcal{G}(1)$. Note that Γ is finite, we can choose m to be the maximum. In this case, as $(\Gamma, X, \mathcal{B}_m) \in \mathcal{G}$, by Lemma 2.3, one of the following three cases occurs:

(5.1) $v_m = k_m = 1$, $\mathcal{B}_m = 1$ and $\Gamma \cong \Pi(\Gamma_{\mathcal{B}_{m-1}}, \Theta_{(m-1,m)})$.

(5.2) $v_m = k_m \ge 2$, and Γ is a multicover of $\Gamma_{\mathcal{B}_m}$.

(5.3) $v_m > k_m \geq 1$, and $(\Gamma, X, \mathcal{B}_m) \in \mathcal{G}$. In this case, let $S_{(m,m+1)} = \{\Gamma_{\mathcal{B}_m}[\tau] | \tau \in V(\Gamma)\}$, $\mathcal{B}_{m+1} = \mathfrak{C}_\Gamma(S_{(m,m+1)})$, then by Lemma 2.3, the $(m+2)$-sequence $\mathcal{B} = \mathcal{B}_0 > \cdots > \mathcal{B}_{m+1}$ satisfies cases **(1)-(4)**, and hence $(\Gamma, X, \mathcal{B}) \in \mathcal{G}(m+1)$, which is contrary to the maximality of m.

Now we are going to bound m. From the analysis above we have $(v,k) \geq v_1 > v_2 > \cdots > v_m \geq 1$ and $v_{i+1} | v_i$ for $i = 0, 1, \cdots, m-1$. So m is no more than the length of the longest factor-chain of $(v,k) = p_1^{n_1} \cdots p_t^{n_t}$, that is, $1 \leq m \leq \sum_{i=1}^t n_i + 1$.

∎

Corollary 2.5 *Let* $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ *such that* $(v,k) = 1$, *then* $\Gamma \cong \Pi(\Gamma_{\mathcal{B}}, \Theta)$, *where* $\Theta = \{(\Gamma_{\mathcal{B}}[\sigma], \Gamma_{\mathcal{B}}[\tau]) | (\sigma, \tau) \in Arc(\Gamma)\}$ *is an* X*-symmetric orbit on* $DSt^r(\Gamma_{\mathcal{B}})$.

3 $\left\llbracket\begin{smallmatrix}\mathbb{r}\\ \mathbb{l}\end{smallmatrix}\right\rrbracket$-double-star graphs

In this section, in order to discuss the higher transitivity of the imprimitive graphs which are not multicovers, we will discuss the $\left\llbracket\begin{smallmatrix}\mathbb{r}\\ \mathbb{l}\end{smallmatrix}\right\rrbracket$-double-stars and the corresponding imprimitive graphs constructed from such double-stars.

In the following three lemmas, we aim to show that in certain conditions, we can obtain a set of $\left\llbracket\begin{smallmatrix}\mathbb{r}\\ \mathbb{l}+1\end{smallmatrix}\right\rrbracket$-double-stars from a set of $\left\llbracket\begin{smallmatrix}\mathbb{r}\\ \mathbb{l}\end{smallmatrix}\right\rrbracket$-double-stars.

Lemma 3.1 *Let* Σ *be a finite* X*-symmetric graph with valency* $\mathbb{b} \geq 2$, Θ *be an* $(X,1)$*-arc-transitive orbit on* $DSt_{\mathbb{l}}^{\mathbb{r}}(\Sigma)$, $S = St(\Theta)$, *where* $\mathbb{l} \geq 1$, $2 \leq \mathbb{r} \leq \mathbb{b}$. *For any* $\mathbb{S} \in S$ *with center* $\sigma_0 \in V(\Sigma)$, *let* $\Theta^+(\mathbb{S}) := \{\mathbb{R} \in S | (\mathbb{S}, \mathbb{R}) \in \Theta\}$, $\Theta^-(\mathbb{S}) := \{\mathbb{L} \in S | (\mathbb{L}, \mathbb{S}) \in \Theta\}$, *and let* $\Theta^*[\mathbb{S}] := \bigcup_{\mathbb{T} \in \Theta^*(\mathbb{S})} \{(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in Arc_{\mathbb{l}+1}(\Sigma, \sigma_0) | (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}, (\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in \mathbb{T}\}$ *for* $* = +, -$. *Then all of the following cases hold:*

(1). $\Theta^*[\mathbb{S}] \subseteq Arc_{\mathbb{l}+1}(\Sigma, \sigma_0)$, $\Theta^*[\mathbb{S}](i) = \mathbb{S}(i)$, *for* $i = 0, 1, \cdots, \mathbb{l}$ *and* $* = +, -$.

(2). $\Theta^*(\mathbb{S})^x = \Theta^*(\mathbb{S}^x)$, $\Theta^*[\mathbb{S}]^x = \Theta^*[\mathbb{S}^x]$ *for any* $x \in X$. *Moreover,* $S^{*2} := \{\Theta^*[\mathbb{S}] | \mathbb{S} \in S\}$ *is* X*-transitive, for* $* = +, -$.

(3). $X_{\mathbb{S}} = X_{\Theta^*[\mathbb{S}]}$, *and if* $\mathbb{S}$ *is* $(X_{\mathbb{S}}, s)$*-arc-transitive, where* s *is an integer with* $1 \leq s \leq \mathbb{l}$, *then* $\Theta^*[\mathbb{S}]$ *is* $(X_{\Theta^*[\mathbb{S}]}, s)$*-arc-transitive. Particularly,* $\Theta^*[\mathbb{S}]$ *is always* $(X_{\Theta^*[\mathbb{S}]}, 1)$*-arc-transitive, for* $* = +, -$.

Proof. The fact that $\mathbb{r} \geq 2$ ensures that $\Theta^*[\mathbb{S}]$ is nonempty.

(1). Clearly, $\Theta^*[\mathbb{S}] \subseteq Arc_{\mathbb{l}+1}(\Sigma, \sigma_0)$, $\Theta^*[\mathbb{S}](\mathbb{l}) \subseteq \mathbb{S}$.

For any $(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}$, and $(\mathbb{L}, \mathbb{R}) \in \Theta$ with $\mathfrak{C}_\Sigma(\mathbb{L}) = \iota$ and $\mathfrak{C}_\Sigma(\mathbb{R}) = \gamma$, as S is X-transitive, there exists some $x \in X$ such that

$\mathbb{L}^x = \mathbb{S}$, then $(\iota^x, \gamma^x) \in \mathbb{S}(1)$. Since S is $(X,1)$-arc-transitive, there exits some $y \in X_{\mathbb{S}}$, such that $(\iota^x, \gamma^x)^y = (\sigma_0, \tau_0)$. Let $\mathbb{T} = \mathbb{R}^{xy}$, then $(\mathbb{S}, \mathbb{T}) = (\mathbb{L}, \mathbb{R})^{xy} \in \Theta$, so $(\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}_{\tau_0} = \mathbb{T}(\mathbb{l}-1)$. Noting $\mathbb{r} \geq 2$, thus there exists some $\tau_{\mathbb{l}} \in V(\Sigma)$ such that $(\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in \mathbb{T}$, and that $(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in Arc_{\mathbb{l}+1}(\Sigma)$. Hence $(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in \Theta^+[\mathbb{S}]$ and so $\mathbb{S} \subseteq \Theta^+[\mathbb{S}](\mathbb{l})$. It follows that $\Theta^+[\mathbb{S}](\mathbb{l}) = \mathbb{S}$.

On the other hand, let x be such that $\mathbb{S} = \mathbb{R}^x$, then similar analysis shows that $\Theta^-[\mathbb{S}](\mathbb{l}) = \mathbb{S}$.

(2). For any $x \in X$, it is not difficult to check that $\Theta^*(\mathbb{S})^x = \Theta^*(\mathbb{S}^x)$. Then $\Theta^*[\mathbb{S}]^x = \bigcup_{\mathbb{T} \in \Theta^*(\mathbb{S})^x} \{(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in Arc_{\mathbb{l}+1}(\Sigma, \sigma_0) | (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}, (\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in \mathbb{T}\} = \bigcup_{\mathbb{T} \in \Theta^*(\mathbb{S}^x)} \{(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in Arc_{\mathbb{l}+1}(\Sigma, \sigma) | (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}, (\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in \mathbb{T}\} = \Theta^*[\mathbb{S}^x]$.

For any $\Theta^*[\mathbb{L}], \Theta^*[\mathbb{R}] \in S^{*2}$, where $\mathbb{L}, \mathbb{R} \in S$, as S is X-transitive, there exists some $x \in X$ such that $\mathbb{L}^x = \mathbb{R}$, then $\Theta^*[\mathbb{L}]^x = \Theta^*[\mathbb{L}^x] = \Theta^*[\mathbb{R}]$. Hence S^{*2} is X-transitive.

(3). It is clear that $X_{\Theta^*[\mathbb{S}]} \leq X_{\Theta^*[\mathbb{S}](\mathbb{l}-1)} = X_{\mathbb{S}}$. For any $x \in X_{\mathbb{S}}$ and $(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}}) \in \Theta^*[\mathbb{S}]$, then $(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}$ and there exists some $\mathbb{T} \in \Theta^*(\mathbb{S})$ such that $(\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}}) \in \mathbb{T}$. As $x \in X_{\mathbb{S}}$, we have $(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}})^x \in Arc_{\mathbb{l}+1}(\Sigma, \sigma_0^x) = Arc_{\mathbb{l}+1}(\Sigma, \sigma_0)$, $(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1})^x \in \mathbb{S}^x = \mathbb{S}$, $(\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}})^x \in \mathbb{T}^x \in \Theta^*[\mathbb{S}^x] = \Theta^*[\mathbb{S}]$. Thus $X_{\mathbb{S}} \leq X_{\Theta^*[\mathbb{S}]}$, and hence $X_{\mathbb{S}} = X_{\Theta^*[\mathbb{S}]}$.

For any $\mathbb{\alpha}, \mathbb{\beta} \in \Theta^*[\mathbb{S}](s) = \mathbb{S}(s)$, as $\mathbb{S}$ is $(X_{\mathbb{S}}, s)$-arc-transitive, there exists some $x \in X_{\mathbb{S}} = X_{\Theta^*[\mathbb{S}]}$ such that $\mathbb{\alpha}^x = \mathbb{\beta}$, hence $\Theta^*[\mathbb{S}]$ is $(X_{\Theta^*[\mathbb{S}]}, s)$-arc-transitive.

∎

Lemma 3.2 *Let Σ be a finite X-symmetric graph with valency $\mathbb{b} \geq 2$, Θ be an $(X,1)$-arc-transitive orbit on $DSt_{\mathbb{l}}^{\mathbb{r}}(\Sigma)$, $S = St(\Theta)$, where $\mathbb{l} \geq 1$, $2 \leq \mathbb{r} \leq \mathbb{b}$. For any $\mathbb{S} \in S$ with center $\sigma_0 \in V(\Sigma)$, we have*

(1). *The following two statements are equal:*

  (1.1). *$\Theta^*[\mathbb{S}] \in St_{\mathbb{l}+1}^{\mathbb{r}}(\Sigma)$ for some $* \in \{+,-\}$.*

  (1.2). *There exists some $\mathbb{T} \in \Theta^{\pm}(\mathbb{S}) := \Theta^+(\mathbb{S}) \cup \Theta^-(\mathbb{S})$ with center τ_0, such that $X_{\mathbb{S}} \cap X_{\tau_0} = X_{\mathbb{T}} \cap X_{\sigma_0}$.*

(2). *If there exists some $\widehat{\mathbb{S}} \in St_{\mathbb{l}+1}^{\mathbb{r}}(\Sigma)$ such that $\widehat{\mathbb{S}}(\mathbb{l}) = \mathbb{S}$, $X_{\widehat{\mathbb{S}}} = X_{\mathbb{S}}$ and for some $(\sigma_0, \gamma_0) \in \widehat{\mathbb{S}}(1)$, $\widehat{\mathbb{S}}_{\gamma_0} \in \Theta^*(\mathbb{S})$, then statements* **(1.1)** *and* **(1.2)** *hold and $\widehat{\mathbb{S}} = \Theta^*[\mathbb{S}]$, for $* = +, -$.*

(3). *If statement* **(1.1)** *or statement* **(1.2)** *holds, then all of the following hold:*

(3.1). *For any* $\mathbb{L}, \mathbb{R} \in S$ *with centers* ι *and* γ*, respectively, such that* $\mathbb{R} \in \Theta^{\pm}(\mathbb{L})$*, then* $X_{\mathbb{L}} \cap X_{\gamma} = X_{\mathbb{R}} \cap X_{\iota}$*.*

(3.2). *For any* $* \in \{+, -\}$*, let* $\sharp = \{+, -\} \backslash \{*\}$*. Let* $\Theta^{*2} = \{(\Theta^*[\mathbb{L}], \Theta^{\sharp}[\mathbb{R}]) | (\mathbb{L}, \mathbb{R}) \in \Theta\}$*. Then for any* $(\widehat{\mathbb{L}}, \widehat{\mathbb{R}}) \in \Theta^{*2}$*,* $X_{\widehat{\mathbb{L}}} \cap X_{\gamma} = X_{\widehat{\mathbb{R}}} \cap X_{\iota}$*, where* ι *and* γ *are the centers of* $\widehat{\mathbb{L}}$ *and* $\widehat{\mathbb{R}}$*, respectively.*

(4). *In case* **(1)***, if we assume further that* Θ *is self-paired, then* $\Theta^+(\mathbb{S}) = \Theta^-(\mathbb{S}) =: \Theta(\mathbb{S})$*,* $\Theta^+[\mathbb{S}] = \Theta^-[\mathbb{S}] =: \Theta[\mathbb{S}]$*,* $S^{+2} = S^{-2} =: S^2$ *and* $\Theta^{+2} = \Theta^{-2} =: \Theta^2$ *is an* X*-symmetric orbit on* $DSt^{\mathbb{r}}_{\mathbb{l}+1}(\Sigma)$*.*

Proof.

(1). Let $\widehat{\mathbb{S}} = \Theta^*[\mathbb{S}]$, $\mathbb{\alpha} = (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \widehat{\mathbb{S}}(\mathbb{l}) = \mathbb{S}$. By the analysis in Lemma 3.1 **(1)**, there exists $\mathbb{T} \in \Theta^*(\mathbb{S})$ with $\mathfrak{C}_{\Sigma}(\mathbb{T}) = \tau_0$. Then $\mathbb{\beta} := (\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{T}(\mathbb{l} - 1)$. As $|\mathbb{T}_{\mathbb{\beta}}| = \mathbb{r} - 1$, there exists exactly $\mathbb{r} - 1$ different vertices $\tau^1_{\mathbb{l}}, \cdots, \tau^{\mathbb{r}-1}_{\mathbb{l}}$ of Σ, such that $(\tau_0, \tau_1, \cdots, \tau^i_{\mathbb{l}}) \in \mathbb{T}$ for $i = 1, \cdots, \mathbb{r} - 1$. Then $(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau^i_{\mathbb{l}}) \in \widehat{\mathbb{S}}$, and hence $|\widehat{\mathbb{S}}_{\mathbb{\alpha}}| \geq \mathbb{r} - 1$.

($\Leftarrow$) If $|\widehat{\mathbb{S}}_{\mathbb{\alpha}}| \geq \mathbb{r}$, then there must be some $\mathbb{R} \in \Theta^*(\mathbb{S}) \setminus \{\mathbb{T}\}$ such that $\mathfrak{C}_{\Sigma}(\mathbb{R}) = \tau_0$. Since Θ is X-transitive, there exists some $x \in X_{\mathbb{S}}$ such that $\mathbb{T}^x = \mathbb{R}$. Then, $x \in X_{\mathbb{S}} \cap X_{\tau_0} \setminus X_{\mathbb{T}} \cap X_{\sigma_0} \neq \emptyset$, a contradiction. Hence $|\widehat{\mathbb{S}}_{\mathbb{\alpha}}| = \mathbb{r} - 1$. It follows from the arbitrariness of $\mathbb{\alpha} \in \widehat{\mathbb{S}}(\mathbb{l})$ and the fact that $\mathbb{S} \in St^{\mathbb{r}}_{\mathbb{l}}(\Sigma)$ that $\widehat{\mathbb{S}} \in St^{\mathbb{r}}_{\mathbb{l}+1}(\Sigma)$.

($\Rightarrow$) If $X_{\mathbb{S}} \cap X_{\tau_0} \neq X_{\mathbb{T}} \cap X_{\sigma_0}$, then there exists some $x \in X_{\mathbb{S}} \cap X_{\tau_0} \setminus X_{\mathbb{T}} \cap X_{\sigma_0}$. Let $\mathbb{R} = \mathbb{T}^x \in \Theta^*(\mathbb{S}^x) = \Theta^*(\mathbb{S})$, then $\mathfrak{C}_{\Sigma}(\mathbb{R}) = \tau_0$, $\mathbb{R} \neq \mathbb{T}$ and $\mathbb{T}(\mathbb{l} - 1) = \mathbb{R}(\mathbb{l} - 1) = \mathbb{S}_{\tau_0}$. Moreover, $\emptyset \neq \mathbb{R} \setminus \mathbb{T} \subseteq \mathbb{R} \setminus \mathbb{R}_{(\tau_0, \sigma_0)}$, that is, there exists some $\mathbb{\tau} := (\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in \mathbb{R} \setminus \mathbb{T}$, such that $\mathbb{\rho} := (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}$ and that $\mathbb{\sigma} := (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau_{\mathbb{l}}) \in \Theta^*[\mathbb{S}]$. By the previous analysis, there exists $\mathbb{r} - 1$ different vertices $\tau^1_{\mathbb{l}}, \tau^2_{\mathbb{l}}, \cdots, \tau^{\mathbb{r}-1}_{\mathbb{l}}$ of Σ such that $(\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau^i_{\mathbb{l}}) \in \mathbb{T}$ and $\mathbb{\sigma}_i := (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}, \tau^i_{\mathbb{l}}) \in \Theta^*[\mathbb{S}] \setminus \{\mathbb{\sigma}\}$, for $i = 1, \cdots, \mathbb{r} - 1$. Then $\{\mathbb{\sigma}, \mathbb{\sigma}_1, \cdots, \mathbb{\sigma}_{\mathbb{r}-1}\} \subseteq \Theta^*[\mathbb{S}]_{\mathbb{\rho}}$, that is, $|\Theta^*[\mathbb{S}]_{\mathbb{\rho}}| \geq \mathbb{r}$, contradicts $S^{*2} \subseteq St^{\mathbb{r}}_{\mathbb{l}+1}(\Sigma)$.

(2). Clearly, $\Theta^*(\mathbb{S}) = \{(\widehat{\mathbb{S}}_{\gamma_0})^x | x \in X_{\widehat{\mathbb{S}}}\} = \{\widehat{\mathbb{S}}_{\gamma^x_0} | x \in X_{\mathbb{S}}\} = \{\widehat{\mathbb{S}}_{\tau_0} | (\sigma_0, \tau_0) \in \widehat{\mathbb{S}}(1)\}$. Thus, $\widehat{\mathbb{S}} = \bigcup_{(\sigma_0, \tau_0) \in \widehat{\mathbb{S}}(1)} \{(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}}) \in Arc_{\mathbb{l}+1}(\Sigma) | (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}, (\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}}) \in \widehat{\mathbb{S}}_{\tau_0}\} = \bigcup_{\mathbb{T} \in \Theta^*(\mathbb{S})} \{(\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}}) \in Arc_{\mathbb{l}+1}(\Sigma) | (\sigma_0, \tau_0, \tau_1, \cdots, \tau_{\mathbb{l}-1}) \in \mathbb{S}, (\tau_0, \tau_1, \cdots, \tau_{\mathbb{l}}) \in \mathbb{T}\} = \Theta^*[\mathbb{S}]$. Then, $\Theta^*[\mathbb{S}] = \widehat{\mathbb{S}} \in St^{\mathbb{r}}_{\mathbb{l}+1}(\Sigma)$, therefore statement **(1.1)** holds, and hence statement **(1.2)** holds.

(3.1). If statement **(1.2)** holds (then statement **(1.1)** holds), as Θ is X-transitive, there exists some $x \in X$ such that $\{\mathbb{S}, \mathbb{T}\}^x = \{\mathbb{L}, \mathbb{R}\}$. Without loss of generality, suppose $\mathbb{S}^x = \mathbb{L}$, then $\mathbb{T}^x = \mathbb{R}$, $\sigma^x = \iota$ and $\tau^x = \gamma$. Hence $X_{\mathbb{L}} \cap X_{\gamma} = X_{\mathbb{S}^x} \cap X_{\tau^x} = (X_{\mathbb{S}} \cap X_{\tau})^x = (X_{\mathbb{T}} \cap X_{\sigma})^x = X_{\mathbb{T}^x} \cap X_{\sigma^x} = X_{\mathbb{R}} \cap X_{\iota}$.

(3.2). By the proof of case **(2)**, $X_{\widehat{\mathbb{L}}} = X_{\mathbb{L}}$ and $X_{\widehat{\mathbb{R}}} = X_{\mathbb{R}}$, thus it follows.

(4). If Θ is self-paired, then $\Theta^{+}(\mathbb{S}) = \{\mathbb{T} \in St(\Theta) | (\mathbb{S}, \mathbb{T}) \in \Theta\} = \{\mathbb{T} \in St(\Theta) | (\mathbb{T}, \mathbb{S}) \in \Theta\} = \Theta^{-}(\mathbb{S})$. Then $S^{+2} = \{\Theta^{+}[\mathbb{T}] | \mathbb{T} \in S\} = \{\Theta^{-}[\mathbb{T}] | \mathbb{T} \in S\}$. Moreover, $\Theta^{+}[\mathbb{S}] = \Theta^{-}[\mathbb{S}]$ and so $\Theta^{+2} = \{(\Theta^{+}(\mathbb{L}), \Theta^{-}(\mathbb{R})) | (\mathbb{L}, \mathbb{R}) \in \Theta\} = \{(\Theta^{-}(\mathbb{L}), \Theta^{+}(\mathbb{R})) | (\mathbb{L}, \mathbb{R}) \in \Theta\} = \Theta^{-2}$.

∎

Lemma 3.3 *Let Σ be a finite X-symmetric graph with valency $\mathbb{b} \geq 2$, Θ be an $(X, 1)$-arc-transitive orbit on $DSt_{\mathbb{l}}^{\mathbb{r}}(\Sigma)$, where $\mathbb{l}$ and $\mathbb{r}$ are integers such that $\mathbb{l} \geq 1$ and $1 \leq \mathbb{r} \leq \mathbb{b}$. Let $\Theta_i = \Theta(i) := \{(\mathbb{L}(i), \mathbb{R}(i)) | (\mathbb{L}, \mathbb{R}) \in \Theta\}$ for $i = 1, \cdots, \mathbb{l}$, then both of the following cases hold:*

(1). *Θ_i is an $(X, 1)$-arc-transitive orbit on $DSt_{i}^{\mathbb{r}}(\Sigma)$ for $i = 1, \cdots, \mathbb{l}$. Moreover, $\Theta_i = \Theta_j(i)$ for $1 \leq i \leq j \leq \mathbb{l}$.*

(2). *If there exists some integer i, where $1 \leq i \leq \mathbb{l}$, such that Θ_i is self-paired, then Θ_j is self-paired for $j = 1, \cdots, i$.*

Proof. Let $\mathbb{S}_i := \mathbb{S}(i)$ for $i = 0, 1, \cdots, \mathbb{l}$.

(1). It is easy to see that $\Theta_i \subset DSt_{i}^{\mathbb{r}}(\Sigma)$. For $(\mathbb{S}(i), \mathbb{T}(i)), (\mathbb{L}(i), \mathbb{R}(i)) \in \Theta_i$, as Θ is X-transitive, there exists some $x \in X$ such that $(\mathbb{S}, \mathbb{T})^x = (\mathbb{L}, \mathbb{R})$, then $(\mathbb{S}(i), \mathbb{T}(i))^x = (\mathbb{L}(i), \mathbb{R}(i))$. For $\mathbb{\alpha}, \mathbb{\beta} \in \mathbb{S}_i(1) = \mathbb{S}(1)$, as Θ is $(X, 1)$-arc-transitive, there exist some $y \in X_{\mathbb{S}} \leq X_{\mathbb{S}_i}$ such that $\mathbb{\alpha}^y = \mathbb{\beta}$. It follows that Θ_i is $(X, 1)$-arc-transitive.

(2). Suppose Θ_i is self-paired, where $1 \leq i \leq \mathbb{l}$, for any $(\mathbb{S}_j, \mathbb{T}_j) \in \Theta_j$, where $1 \leq j \leq i$, there exists $(\mathbb{S}_i, \mathbb{T}_i) \in \Theta$ such that $\mathbb{S}_j = \mathbb{S}_i(j), \mathbb{T}_j = \mathbb{T}_i(j)$. As $(\mathbb{T}_i, \mathbb{S}_i) \in \Theta_i$, hence $(\mathbb{T}_j, \mathbb{S}_j) = (\mathbb{T}_i(j), \mathbb{S}_i(j)) \in \Theta_j$, it follows that Θ_j is self-paired.

∎

Let Σ be a finite X-symmetric graph, Θ be an $(X, 1)$-arc-transitive orbit on $DSt_{\mathbb{l}}^{\mathbb{r}}(\Sigma)$, $\mathbb{S} \in St(\Theta)$. The following theorem shows that the subgroup sequence $X_{\mathbb{S}(i)}$, for $i = 0, 1, \cdots$, decreases strictly.

Theorem 3.4 *Let Σ be a finite X-symmetric graph with valency $\mathbb{b} \geq 3$, Θ be an $(X, 1)$-arc-transitive orbit on $DSt_{\mathbb{l}}^{\mathbb{r}}(\Sigma)$, where $\mathbb{l}$ and $\mathbb{r}$ are integers such that $\mathbb{l} \geq 1$ and $2 \leq \mathbb{r} \leq \mathbb{b}$. Then there exists some integer $\mathbb{h}$ with $1 \leq \mathbb{h} \leq \mathbb{l}$ such that for any $(\mathbb{S}, \mathbb{T}) \in \Theta$ with centers $\sigma, \tau \in V(\Sigma)$, respectively, $X_\sigma = X_{\mathbb{S}(0)} > X_{\mathbb{S}(1)} > \cdots > X_{\mathbb{S}(\mathbb{h})} = \cdots = X_{\mathbb{S}(\mathbb{l})} = X_{\mathbb{S}} > 1$. Moreover, either of the following two cases occurs:*

(1). *$\mathbb{h} = \mathbb{l}$, and $X_{\mathbb{S}(i)} \cap X_\tau \neq X_{\mathbb{T}(i)} \cap X_\sigma$, for $i = 1, \cdots, \mathbb{l}$.*

(2). $\mathbb{h} \leq \mathbb{l}$, *there exists a series* $\{(\mathbb{S}_i, \mathbb{T}_i) \in \Theta_i \in DSt_i^{\mathbb{r}}(\Sigma) | i = 1, 2, \cdots\}$, *such that all of the following hold:*

  (2.1). $(\mathbb{S}_i, \mathbb{T}_i) = (\mathbb{S}(i), \mathbb{T}(i))$ *and* $\Theta_i = \Theta(i)$ *for* $i = 1, 2, \cdots, \mathbb{l}$.

  (2.2). $\mathbb{S}_{i+1} = \Theta_i^+[\mathbb{S}_i]$, $\mathbb{T}_{i+1} = \Theta_i^-[\mathbb{T}_i] \in St_{i+1}^{\mathbb{r}}(\Sigma)$, $\Theta_{i+1} = \Theta_i^2$, *for* $i = h, h+1, \cdots$.

  (2.3). $X_\sigma > X_{\mathbb{S}_1} > \cdots > X_{\mathbb{S}_{\mathbb{h}}} = X_{\mathbb{S}_{\mathbb{h}+1}} = \cdots$. *Moreover,* $X_{\mathbb{S}(i)} \cap X_\tau \neq X_{\mathbb{T}(i)} \cap X_\sigma$, *for* $i = 1, \cdots, \mathbb{h} - 1$, *and* $X_{\mathbb{S}_i} \cap X_\tau = X_{\mathbb{T}_i} \cap X_\sigma$, *for* $i = h, h+1, \cdots$.

Proof. By Lema 3.2, $X_{\mathbb{S}(i)} = X_{\mathbb{S}(i+1)}$ for some integer i with $1 \leq i \leq \mathbb{l} - 1$, if and only if $X_{\mathbb{S}(i)} \cap X_\tau = X_{\mathbb{T}(i)} \cap X_\sigma$. Thus if $X_{\mathbb{S}(i)} \cap X_\tau \neq X_{\mathbb{T}(i)} \cap X_\sigma$ for all $1 \leq i \leq \mathbb{l}$, then **(1)** holds.

Otherwise, choose $\mathbb{h}$ to be the minimal such that $X_{\mathbb{S}(\mathbb{h})} \cap X_\tau = X_{\mathbb{T}(\mathbb{h})} \cap X_\sigma$, where $1 \leq \mathbb{h} \leq \mathbb{l}$. Let $\mathbb{S}_i = X_{\mathbb{S}(i)}$, $\Theta_i = \Theta(i)$, for $i = 1, \cdots, \mathbb{h}$. For any $i \geq \mathbb{h}$ such that $X_{\mathbb{S}_i} \cap X_\tau = X_{\mathbb{T}_i} \cap X_\sigma$, let $\mathbb{S}_{i+1} = \Theta_i^+[\mathbb{S}_i]$, $\mathbb{T}_{i+1} = \Theta_i^-[\mathbb{T}_i]$, then by Lema 3.2, $X_{\mathbb{S}_{i+1}} \cap X_\tau = X_{\mathbb{T}_{i+1}} \cap X_\sigma$, and hence it follows.

∎

Hereafter, for $(\Sigma, X, \Theta) \in \mathbb{G}$, let $\hat{p}(\Sigma, X, \Theta) := (\mathbb{b}, \mathbb{r}, \mathbb{l}, \mathbb{h})$.

Construction 3.5 *Let* Σ *be a finite* X*-symmetric graph with valency* $\mathbb{b} \geq 2$, Θ *be an* X*-symmetric orbit on* $DSt_{\mathbb{l}}^{\mathbb{r}}(\Sigma)$, *where* $\mathbb{b}, \mathbb{r}$ *and* $\mathbb{l}$ *are integers such that* $\mathbb{l} \geq 1$ *and* $2 \leq \mathbb{r} \leq \mathbb{b}$. *Then the* $\llbracket {\mathbb{r} \atop \mathbb{l}} \rrbracket$*-double-star graph* $\Pi(\Sigma, \Theta)$ *with respect to* Θ *is defined to be the graph with vertex set* $St(\Theta) := \{\mathbb{L}, \mathbb{R} | (\mathbb{L}, \mathbb{R}) \in \Theta\}$ *and edge set* $\{\{\mathbb{L}, \mathbb{R}\} | (\mathbb{L}, \mathbb{R}) \in \Theta\}$.

For any finite graph Ψ, denote by $Com(\Psi)$ the set of the connected components of Ψ, and let $c(\Psi) := |Com(\Psi)|$. Let Γ be a finite X-symmetric graph with an X-invariant partition $\mathcal{B}$ on $V(\Gamma)$ such that $\Gamma_{\mathcal{B}}$ is nonempty. Then $c := c(\Gamma[B, C])$ is independent of the choice of $(B, C) \in Arc(\Gamma_{\mathcal{B}})$. Let h be the maximal integer such that $1 \leq h \leq m$ and $d = d_{h-1}$. Let $\hat{p}(\Gamma, X, \mathcal{B}) := (v, k, r, b, d, m, c, h)$.

Let Σ be a finite X-symmetric graph. In the theorem below, we study the structure of the $\llbracket {\mathbb{r} \atop \mathbb{l}} \rrbracket$-double-star graphs of Σ by counting some useful parameters. As a byproduct, we give a sufficient condition for the existence of $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ such that $\Gamma_{\mathcal{B}} \cong \Sigma$ and Γ is (X, s)-arc-transitive, for some inter s with $1 \leq s \leq \mathbb{l}$.

Theorem 3.6 *Let* $(\Sigma, X, \Theta) \in \mathbb{G}$ *with* $\hat{p}(\Sigma, X, \Theta) := (\mathbb{b}, \mathbb{r}, \mathbb{l}, \mathbb{h})$, $\Pi = \Pi(\Sigma, \Theta)$ *and* $S = St(\Theta)$. *Then* $(\Pi, X, \mathcal{S}) \in \mathcal{G}$ *and all of the following hold:*

(1). *For any* $\mathbb{S} \in S$, *let* $S_{\mathbb{S}(i)} := \{\mathbb{T} \in S | \mathbb{T}(i) = \mathbb{S}(i)\}$, $\mathcal{S}_i := \{S_{\mathbb{S}(i)} | \mathbb{S} \in S\}$ *for* $i = 1, \cdots, \mathbb{h}$. *Then* $\mathcal{S}_i$ *is an* X*-invariant partition on* $S = V(\Pi)$, *for* $i = 1, \cdots, \mathbb{h}$, *such that* $\mathcal{S} = \mathcal{S}_0 > \mathcal{S}_1 > \cdots > \mathcal{S}_{\mathbb{h}} = 1$.

(2). $\Pi_i := \Pi(\Sigma, \Theta(i)) \cong \Pi_{\mathcal{S}_i}$ *for* $i = 1, \cdots, \mathbb{h}$. *Moreover, for any* $\mathbb{S} \in S$, *let* $\mathbb{S}_{(i,i+1)} := \{(\mathbb{S}(i), \mathbb{T}(i)) | \mathbb{T} \in \Theta(\mathbb{S})\}$, $\Theta_{(i,i+1)} := \{(\mathbb{L}_{(i,i+1)}, \mathbb{R}_{(i,i+1)}) | (\mathbb{L}, \mathbb{R}) \in \Theta\}$, *then* $\mathbb{S}_{(i,i+1)}$ *is an* $X_{\mathbb{S}(i+1)}$*-symmetric* $\mathbb{r}$*-star of* Π_i *while* $\Theta_{(i,i+1)}$ *is an* X*-symmetric orbit on* $DSt^{\mathbb{r}}(\Pi_i)$ *such that* $\Pi_{i+1} \cong \Pi(\Pi_i, \Theta_{(i,i+1)})$, *for* $i = 1, \cdots, \mathbb{h} - 1$.

(3). *For any* $(\mathbb{S}, \mathbb{T}) \in \Theta$, *we have* $Com(\Gamma[S_{\mathbb{S}(0)}, S_{\mathbb{T}(0)}]) \supset Com(\Gamma[S_{\mathbb{S}(1)}, S_{\mathbb{T}(1)}]) \supset \cdots \supset Com(\Gamma[S_{\mathbb{S}(\mathbb{h}-1)}, S_{\mathbb{T}(\mathbb{h}-1)}])$.

(4). *Let* $\hat{p}(\Pi_i, X, \mathcal{S}_i) = (v_i, k_i, r_i, b_i, d_i, m_i, c_i, h_i)$, *then* $m_i = i$, $h_i = i$, $r_i = \mathbb{r}$, $d_i = val(\Pi)/\mathbb{r}$ *for* $i = 1, 2, \cdots, \mathbb{h}$. *Additionally,* $k/c = k_0/c_0 = k_1/c_1 = \cdots = k_{\mathbb{h}-1}/c_{\mathbb{h}-1} \geq d$.

(5). *For any* $(\mathbb{S}, \mathbb{T}) \in \Theta$, *where* $\mathfrak{C}_\Sigma(\mathbb{S}) = \sigma$ *and* $\mathfrak{C}_\Sigma(\mathbb{T}) = \tau$, *either of the following cases occurs:*

(5.1). $d \geq 2$, $\mathbb{h} = \mathbb{l}$ *and* $X_{\mathbb{S}} \cap X_\tau \neq X_{\mathbb{T}} \cap X_\sigma$. Π *is* $(X, 1)$*-transitive.*

(5.2). $d = 1$, $\mathbb{h} \leq \mathbb{l}$, *and* $X_{\mathbb{S}} \cap X_\tau = X_{\mathbb{T}} \cap X_\sigma$. *Moreover, if* $\mathbb{S}$ *is* $(X_{\mathbb{S}}, s)$*-arc transitive for some integer* $1 \leq s \leq \mathbb{l}$, *then* Π *is* (X, s)*-arc-transitive.*

Proof.

(1). For any integer i with $1 \leq i \leq \mathbb{h}$, clearly, $\mathcal{S}_i$ is a partition of $\mathcal{S} = V(\Pi)$. For any $\mathbb{S} \in S$ and $x \in X$, $S_{\mathbb{S}(i)}^x = \{\mathbb{T} | \mathbb{T}(i) = \mathbb{S}(i)\}^x = \{\mathbb{T} \in S | \mathbb{T}(i) = \mathbb{S}(i)^x\} = S_{\mathbb{S}(i)^x}$. Hence $\mathcal{S}_i$ is an X-invariant partition. Moreover, it is not difficult to check that $X_{S_{\mathbb{S}(i)}} = X_{\mathbb{S}(i)}$ for $i = 1, \cdots, \mathbb{h}$.

For any integer i with $1 \leq i \leq \mathbb{h} - 1$ and $\mathbb{T} \in S_{\mathbb{S}(i+1)}$, then $\mathbb{T}(i+1) = \mathbb{S}(i+1)$ and hence $\mathbb{T}(i) = \mathbb{S}(i)$, that is, $\mathbb{T} \in S_{\mathbb{S}(i)}$ and so $S_{\mathbb{S}(i+1)} \subseteq S_{\mathbb{S}(i)}$. If $S_{\mathbb{S}(i+1)} = S_{\mathbb{S}(i)}$, then $X_{\mathbb{S}(i+1)} = X_{S_{\mathbb{S}(i+1)}} = X_{S_{\mathbb{S}(i)}} = X_{\mathbb{S}(i)}$. Then $\mathbb{h} \leq i$, a contradiction. Thus $S_{\mathbb{S}(i+1)} \subset S_{\mathbb{S}(i)}$. It follows that $\mathcal{S} = \mathcal{S}_1 > \cdots > \mathcal{S}_{\mathbb{h}}$. Note that $S_{\mathbb{S}(\mathbb{h})} = \{\mathbb{S}\}$ for any $\mathbb{S} \in S$, hence $\mathcal{S}_{\mathbb{h}} = 1$.

(2). For any $(\mathbb{S}, \mathbb{T}) \in \Theta$, as Θ is self-paired, there exists some $z \in X$ such that $(\mathbb{S}, \mathbb{T})^z = (\mathbb{T}, \mathbb{S})$, then $(\mathbb{S}(i), \mathbb{T}(i))^z = (\mathbb{T}(i), \mathbb{S}(i)) \in Arc(\Pi_i)$, $(S_{\mathbb{S}(i)}, S_{\mathbb{T}(i)})^z = (S_{\mathbb{T}(i)}, S_{\mathbb{S}(i)}) \in Arc(\Pi_{\mathcal{S}_i})$. Let $G = X_{\mathbb{S}(i)}$ then $G = X_{S_{\mathbb{S}(i)}}$. Thus $\Pi_i \cong Cos(X, G, GzG) \cong \Pi_{\mathcal{S}_i}$.

For any $\tau \in V(\Sigma)$ such that $(\sigma, \tau) \in \mathbb{S}(1)$, we have showed that there exists some $\mathbb{T} \in S$ with center τ such that $(\mathbb{S}, \mathbb{T}) \in \Theta$, it follows that $|\mathbb{S}_{(i,i+1)}| \geq \mathbb{r}$. On the other hand, if there exists some $\mathbb{R}$ with center τ such that $(\mathbb{S}, \mathbb{R}) \in \Theta$, then $\mathbb{R}(i) = \mathbb{T}(i) = \{(\tau, \tau_1, \cdots, \tau_i) \in Arc_i(\Sigma) | (\sigma, \tau, \tau_1, \cdots, \tau_i) \in \mathbb{S}(i+1)\} \cup \{(\tau, \sigma, \sigma_1, \cdots, \sigma_{i-1}) \in Arc_i(\Sigma) | (\sigma, \sigma_1, \cdots, \sigma_{i-1}) \in \mathbb{S}(i-1)\}$. It follows that $|\mathbb{S}_{(i,i+1)}| = \mathbb{r}$. It is not difficult to check that $X_{\mathbb{S}_{(i,i+1)}} = X_{\mathbb{S}(i+1)}$ and $\mathbb{S}_{(i,i+1)} \in St^{\mathbb{r}}(\Pi_i)$ is $X_{\mathbb{S}_{(i,i+1)}}$-symmetric.

For any $z \in X$ such that $(\mathbb{S}, \mathbb{T})^z = (\mathbb{T}, \mathbb{S})$, clearly, $(\mathbb{S}_{(i,i+1)}, \mathbb{T}_{(i,i+1)})^z = (\mathbb{T}_{(i,i+1)}, \mathbb{S}_{(i,i+1)}) \in \Theta_{(i,i+1)}$. It follows that $\Theta_{(i,i+1)}$ is a self-paired X-symmetric orbit on $DSt^{\mathbbm{r}}(\Pi_i)$. Let $G = X_{\mathbb{S}_{(i,i+1)}} = X_{\mathbb{S}(i+1)}$, then $\Pi_{i+1} \cong Cos(X, G, GzG) \cong \Pi(\Pi_i, \Theta_{(i,i+1)})$.

(3). Let $\sigma = \mathfrak{C}_\Sigma(\mathbb{S})$, $\tau = \mathfrak{C}_\Sigma(\mathbb{T})$. For any $\mathbb{L} \in S_\sigma \setminus \{\mathbb{S}\}$, such that $\mathbb{S}$ and $\mathbb{L}$ are in the same connected component of $\Pi[S_\sigma, S_\tau]$, then there exists a $2j$-path $[\mathbb{S} = \mathbb{S}_0, \mathbb{T}_1, \mathbb{S}_1, \cdots, \mathbb{T}_j, \mathbb{S}_j = \mathbb{L}]$ of $\Pi[S_\sigma, S_\tau]$), where $j \geq 1$ and $\mathbb{S}_i \in S_\sigma, \mathbb{T}_i \in S_\tau$ for $i = 0, 1, \cdots, j$. For any $0 \leq i \leq j-1$, as $(\mathbb{T}_{i+1}, \mathbb{S}_i), (\mathbb{T}_{i+1}, \mathbb{S}_{i+1}) \in \Theta$, hence $\mathbb{S}_i(\mathbbm{h}-1) = \mathbb{S}_{i+1}(\mathbbm{h}-1)$. And so $\mathbb{L}(\mathbbm{h}-1) = \mathbb{S}(\mathbbm{h}-1) \in S_{\mathbb{S}(\mathbbm{h}-1)} \in \mathcal{S}_{\mathbbm{h}-1}$. Thus it follows.

(4). By the analysis in **(1)** and **(2)**, we have that $m_i = i$, $h_i = i$, $r_i = \mathbbm{r}$, $d_i = val(\Pi)/\mathbbm{r}$, for $i = 1, 2, \cdots, \mathbbm{h}$. By **(3)** we have $k_i/c_i = k/c \geq d$ for $i = 1, 2, \cdots, \mathbbm{h}-1$.

(5.1). If $X_{\mathbb{S}} \cap X_\tau \neq X_{\mathbb{T}} \cap X_\sigma$, then there exists $x \in X_{\mathbb{S}} \cap X_\tau \setminus X_{\mathbb{T}} \cap X_\sigma \neq \emptyset$. Then $(\mathbb{S}, \mathbb{T}), (\mathbb{S}, \mathbb{T}^x) \in Arc(\Pi[S_\sigma, S_\tau])$, where $\mathbb{T} \neq \mathbb{T}^x$. It follows that $d \geq 2$. As $r = \mathbbm{r} \geq 2$, Π is $(X, 1)$-transitive.

(5.2). Otherwise, $X_{\mathbb{S}} \cap X_\tau = X_{\mathbb{T}} \cap X_\sigma$. If there exits $\mathbb{R} \in S_\tau$ such that $(\mathbb{S}, \mathbb{R}) \in \Theta$, then there exits some $x \in X$ such that $(\mathbb{S}, \mathbb{T})^x = (\mathbb{S}, \mathbb{R})$. Then $x \in X_{\mathbb{S}} \cap X_\tau = X_{\mathbb{T}} \cap X_\sigma$, and hence $\mathbb{R} = \mathbb{T}^x$. It follows that $d = 1$.

If $\mathbb{S}$ is $(X_{\mathbb{S}}, s)$-arc-transitive, for some integer s with $1 \leq s \leq \mathbbm{l}$, let $(\mathbb{S} = \mathbb{S}_0, \mathbb{S}_1, \cdots, \mathbb{S}_s), (\mathbb{S} = \mathbb{L}_0, \mathbb{L}_1, \cdots, \mathbb{L}_s) \in Arc_s(\Pi, \mathbb{S})$, and let $\mathfrak{C}_\Sigma(\mathbb{S}_i) = \sigma_i$, $\mathfrak{C}_\Sigma(\mathbb{L}_i) = \iota_i$ for $i = 0, 1, \cdots, s$. Clearly $\mathbbm{\sigma}_i := (\sigma_{s-i}, \sigma_{s-i+1}, \cdots, \sigma_s) \in Arc_i(\Sigma)$ for $i = 1, \cdots, s$, and $\mathbbm{\sigma}_1 \in \mathbb{S}_{s-1}(1)$. Suppose $\mathbbm{\sigma}_i \in \mathbb{S}_{s-i}(i)$ for some $1 \leq i \leq s-1$, then as $(\mathbb{S}_{s-i-1}, \mathbb{S}_{s-i}) \in \Theta$ and $\sigma_{s-i-1} \neq \sigma_{s-i+1}$, we have $\mathbbm{\sigma}_{i+1} \in \mathbb{S}_{s-i-1}(i+1)$. Hence we have $\mathbbm{\sigma}_s \in \mathbb{S}(s)$. Similarly, $\mathbbm{\beta}_s := (\iota_0, \cdots, \iota_s) \in \mathbb{S}(s)$. Then there exists some $x \in \mathbb{S}$ such that $\mathbbm{\alpha}_s = \mathbbm{\beta}_s^x$. If there exists some $0 \leq i \leq s-1$ such that $\mathbb{S}_{i+1} \neq \mathbb{L}_{i+1}^x$, choose i to be the minimal. Then $(\mathbb{S}_i, \mathbb{S}_{i+1}), (\mathbb{S}_i, \mathbb{L}_{i+1}^x) \in Arc(\Pi[\sigma_i, \sigma_{i+1}])$, that is, $d \geq 2$, a contradiction. Hence $(\mathbb{S}_0, \mathbb{S}_1, \cdots, \mathbb{S}_s) = (\mathbb{L}_0, \mathbb{L}_1, \cdots, \mathbb{L}_s)^x$, it follows that Π is (X, s)-arc-transitive.

∎

Here we give some examples to show the construction of (X, s)-arc-transitive imprimitive graphs which are not multicovers.

Example 3.7 Let Σ *be the complete graph with vertex set* $\{1, 2, 3, 4, 5\}$, $X = A_5 < S_5 = Aut(\Sigma)$, $z = (15)(24) \in X$, *then* Σ *is* $(X, 2)$*-arc-regular. Let* $\mathbb{S} = \{(1, 5), (1, 4), (1, 3)\}$, $\mathbb{T} = \{(5, 1), (5, 2), (5, 3)\}$, $\Theta = \{(\mathbb{S}, \mathbb{T})^x | x \in X\}$. *It is not difficult to check that* $(\mathbb{S}, \mathbb{T})^z = (\mathbb{T}, \mathbb{S})$ *and* Θ *is an* X*-symmetric orbit on* $DSt^3(\Sigma)$ *such that* $X_{\mathbb{S}} \cap X_5 = X_{\mathbb{T}} \cap X_1 = 1$. *Let* $\Gamma = \Pi(\Sigma, \Theta)$, *then* Γ *is a connected cubic* $(X, 1)$*-arc-regular graph.*

For any integer $n \geq 1$ and $1 \leq q \leq n$, let $[n] := \{1, 2, \cdots, n\}$, and let $[n]^{(q)}$ be the set of the q-subsets of $[n]$. For $n \geq 2$, the odd graph O_n is defined to be the graph with vertex set $[2n-1]^{(n-1)}$, and two vertices are adjacent of and only if the corresponding subsets are disjoint. By [1], $Aut(O_n) = Sym([2n-1])$.

Example 3.8 Let Σ *be the Peterson graph* O_3*, and let* $X = A_5 < S_5 = Aut(O_3)$*, then* Σ *is* $(X, 2)$*-arc-regular. Let* $x = (13524) \in X$*. On the one hand, let* $\sigma = \{1, 2\} \in V(\Sigma)$*. For any integer* $i \geq 0$*, let* $\mathbb{S}_i = \{(\sigma, \sigma^y, \sigma^{y^2}, \cdots, \sigma^{y^i}) | y \in \{x, x^{-1}\}\}$*, then* $\mathbb{S}_i \in St_i^2(\Sigma)$ *with center* σ*, for* $i \geq 1$*. Moreover,* $X_\sigma = X_{\mathbb{S}_0} > X_{\mathbb{S}_1} = X_{\mathbb{S}_2} = \cdots$*, where* $X_\sigma = \{(1), (345), (543), (12)(34), (12)(35), (12)(45)\}$*,* $X_{\mathbb{S}_i} = \{(1), (12)(35)\}$ *and* $\mathbb{S}_i$ *is* $(X_{\mathbb{S}_i}, i)$*-arc-transitive for all* $i \geq 1$*.*

On the other hand, let $\tau = \{3, 4\}$*, and for* $i \geq 0$*, let* $\mathbb{T}_i = \{(\tau, \tau^y, \tau^{y^2}, \cdots, \tau^{y^i}) | y \in \{x, x^{-1}\}\}$*. Then* $\mathbb{T}_i \in St_i^2(\Sigma)$ *with center* τ*, for* $i = 1, 2, \cdots$*. Moreover,* $X_\tau = X_{\mathbb{T}_0} > X_{\mathbb{T}_1} = X_{\mathbb{T}_2} = \cdots$*, where* $X_\tau = \{(1), (125), (521), (34)(12), (34)(15), (34)(25)\}$*,* $X_{\mathbb{T}_i} = \{(1), (34)(25)\}$ *and* $\mathbb{T}_i$ *is* $(X_{\mathbb{T}_i}, i)$*-arc-transitive for all* $i \geq 1$*. Additionally, it is not difficult to check that* $\Theta_i := \{(\mathbb{S}_i, \mathbb{T}_i)^x | x \in X\}$ *is a self-paired* (X, i)*-arc-transitive orbit on* $DSt_i^2(\Sigma)$ *for all* $i \geq 1$*. Let* $\Pi = \Pi(\Sigma, \Theta)$*, and let* $\Psi = [\{1, 2\}, \{3, 4\}, \{5, 1\}, \{2, 3\}, \{4, 5\}, \{1, 2\}]$*, then* Ψ *is a* 5*-cycle of* Σ *such that* $\Pi \cong 6\Psi$ *is* (X, s)*-arc-transitive for all* $s \geq 1$*.*

Example 3.9 Consider the odd graph O_4*, let* $X = A_7 < S_7 = Aut(O_4)$*, then* O_4 *is* $(X, 3)$*-transitive (see, for example, [3]). Set*

$$\mathbb{S} := \left\{ \begin{array}{c} (123, 456, 127) \\ (123, 456, 137) \\ (123, 457, 126) \\ (123, 457, 136) \\ (123, 567, 124) \\ (123, 567, 134) \end{array} \right\}; \mathbb{T} := \left\{ \begin{array}{c} (456, 123, 457) \\ (456, 123, 567) \\ (456, 127, 345) \\ (456, 123, 356) \\ (456, 137, 245) \\ (456, 137, 256) \end{array} \right\}.$$

Clearly, $\mathbb{S} \in St_2^3(O_4)$*, and simple counting shows that* $X_{\mathbb{S}} = (Sym(\{4, 6, 7\}) \times Sym(\{2, 3\})) \cap A_7 = \{(1), (467), (764), (46)(23), (47)(23), (67)(23)\}$ *and that* $\mathbb{S}$ *is* $(X_{\mathbb{S}}, 2)$*-arc-transitive. One can check that* $(\mathbb{S}, \mathbb{T})$ *is a* $\left[\!\left[{3 \atop 2}\right]\!\right]$*-double-star of* O_4 *with* $(\mathbb{S}, \mathbb{T})^z = (\mathbb{T}, \mathbb{S})$*, that is,* $\Theta := \{(\mathbb{S}, \mathbb{T})^x | x \in X\}$ *is a self-paired* $(X, 2)$*-arc-transitive orbit on* $DSt_2^3(O_4)$*. Moreover,* $X_{\mathbb{S}} \cap X_{\{4,5,6\}} = X_{\mathbb{T}} \cap X_{\{1,2,3\}} = \{(1), (23)(46)\}$*.*

Let $\Gamma = O_3$*,* $\widetilde{\Gamma} = \widetilde{\Gamma}(K, \phi)$ *be the double covering graph of* O_3*, where* $K = Z_2 = \{1, a\}$ *and the function* ϕ*, which assigns* a *to each arc of* O_3 *is a* Z_2*-chain of* O_3*. Let* $\Pi = \Pi(O_4, \Theta)$*, then simple analysis shows that* $\Pi \cong 12\widetilde{\Gamma}$ *is* $(X, 2)$*-transitive. In the Figure 2, we depict* O_4 *and one of the connected components* $\widetilde{\Gamma}$ *(up to isomorphism) of* Π*.*



the odd graph O_4

a 2-fold cover of the peterson graph

Figure 2. An $(X, 2)$-arc-transitive graph with an $(X, 3)$-arc-transitive quotient

Example 3.10 For any integer $n \geq 3$, let Σ be the complete bipartite graph with bipartition $\{\iota_i | i = 1, 2, \cdots, n\} \cup \{\gamma_i | i = 1, 2, \cdots, n\}$, $X = Aut(\Sigma) = Sym([n]) \wr Sym(\{\iota, \gamma\})$, $\mathbb{S} = \{(\iota_1, \gamma_i, \iota_j) | i \in [n] \setminus \{1\}, j \in [n] \setminus \{1, i\}\}$, $\mathbb{T} = \{(\gamma_2, \iota_i, \gamma_j) | i \in [n] \setminus \{2\}, j \in [n] \setminus \{2, i\}\}$, then $(\mathbb{S}, \mathbb{T}) \in DSt_2^{n-1}(\Sigma)$, $X_{\mathbb{S}} = Sym([n] \setminus \{1\})$ and $\mathbb{S}$ is $(X_{\mathbb{S}}, 2)$-arc-transitive. Let $z = (\iota_1\ \gamma_2)(\iota_2\ \gamma_1)(\iota_3\ \gamma_3) \cdots (\iota_i\ \gamma_i)(\iota_n\ \gamma_n) \in X$, then $(\mathbb{S}, \mathbb{T})^z = (\mathbb{T}, \mathbb{S})$, that is, $\Theta := \{(\mathbb{S}, \mathbb{T})^x | x \in X\}$ is a self-paired $(X, 2)$-arc-transitive orbit on $DSt_2^{n-1}(\Sigma)$. Moreover, it is not difficult to check that $X_{\mathbb{S}} \cap X_{\iota_2} = X_{\mathbb{T}} \cap X_{\gamma_1} = Sym([n] \setminus \{1, 2\})$. Let $\Pi = \Pi(\Sigma, \Theta)$, then $\Pi \cong n(K_{n,n} - nK_2)$ is $(X, 2)$-arc-transitive.

In Section 2, for $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, we have defined $S_i := \{\Gamma_{\mathcal{B}_i}[\tau] | \tau \in V(\Gamma)\}$, $\Theta_i := \{(\Gamma_{\mathcal{B}_i}[\tau], \Gamma_{\mathcal{B}_i}[\sigma]) | (\tau, \sigma) \in Arc(\Gamma)\}$, $\mathcal{B}_{i+1} = \mathfrak{C}_\Gamma(S_i)$ for $i = 0, 1, \ldots, m-1$. For the sake of convenient, we continue to define $S_i := \{\Gamma_{\mathcal{B}_i}[\tau] | \tau \in V(\Gamma)\}$, $\Theta_i := \{(\Gamma_{\mathcal{B}_i}[\tau], \Gamma_{\mathcal{B}_i}[\sigma]) | (\tau, \sigma) \in Arc(\Gamma)\}$, $\mathcal{B}_{i+1} = \mathfrak{C}_\Gamma(S_i)$ for $i = m, m+1, \ldots$. Then it is clear that $S_m = S_{m+1} = \cdots$, $\Theta_m = \Theta_{m+1} = \cdots$, $\mathcal{B}_m = \mathcal{B}_{m+1} = \cdots$, and $\Gamma_{\mathcal{B}_m} = \Gamma_{\mathcal{B}_{m+1}} = \cdots$.

Let Γ be a finite graph with a partition $\mathcal{B}$ on $V(\Gamma)$. For any $\sigma \in V(\Gamma)$, let $\mathcal{B}(\sigma)$ be the block of $\mathcal{B}$ which contains σ. Let $Arc_i(\Gamma_{\mathcal{B}}, \sigma) := \{(\mathcal{B}(\sigma), \mathcal{B}(\sigma_1), \cdots, \mathcal{B}(\sigma_i)) \in Arc_i(\Gamma_{\mathcal{B}}) | (\sigma, \sigma_1, \cdots, \sigma_i) \in Arc_i(\Gamma)\}$. Then $Arc_i(\Gamma_{\mathcal{B}}, \sigma) \subseteq Arc_i(\Gamma_{\mathcal{B}}, \mathcal{B}(\sigma))$ is X-symmetric.

In order to complete the proof of our main result, we need to give some necessary lemmas and theorems at the end of this section.

Lemma 3.11 *Let $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, $\sigma, \iota \in V(\Gamma)$, i, j be two nonnegative integers, then $Arc_i(\Gamma_{\mathcal{B}_j}, \sigma) = Arc_i(\Gamma_{\mathcal{B}_j}, \iota)$ if and only if $\mathcal{B}_{i+j}(\sigma) = \mathcal{B}_{i+j}(\iota)$. Moreover, $X_{Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)} = X_{\mathcal{B}_{i+j}(\sigma)}$.*

Proof. Fix j, then the case of $i = 0$ is trivial.

($\Leftarrow$) Suppose it follows for some $i \geq 0$. For any $(\sigma, \sigma_1, \cdots, \sigma_i, \sigma_{i+1}) \in Arc_{i+1}(\Gamma)$ such that $(\mathcal{B}_j(\sigma), \mathcal{B}_j(\sigma_1), \cdots, \mathcal{B}_j(\sigma_{i+1})) \in Arc_{i+1}(\Gamma_{\mathcal{B}_j})$, as $\iota \in \mathcal{B}_{i+1}(\sigma)$, there exists $\iota_1 \in \Gamma(\iota)$ such that $\iota_1 \in \mathcal{B}_i(\sigma)$, then $Arc_i(\Gamma_{\mathcal{B}_j}(\sigma_1)) = Arc_i(\Gamma_{\mathcal{B}_j}(\sigma_1))$, hence there exists $(\iota_1, \cdots, \iota_{i+1}) \in Arc_i(\Gamma)$ such that $(\mathcal{B}_j(\iota_1), \cdots, \mathcal{B}_j(\iota_{i+1})) \in Arc_i(\Gamma_{\mathcal{B}_j})$. Then $(\iota, \iota_1, \cdots, \iota_{i+1}) \in Arc_{i+1}(\Gamma)$ and $(\mathcal{B}_j(\iota), \mathcal{B}_j(\iota_1), \cdots, \mathcal{B}_j(\iota_{i+1})) \in Arc_{i+1}(\Gamma_{\mathcal{B}_j})$. Thus $Arc_{i+1}(\Gamma_{\mathcal{B}_j}, \sigma) \subseteq Arc_{i+1}(\Gamma_{\mathcal{B}_j}, \iota)$. Similarly, $Arc_{i+1}(\Gamma_{\mathcal{B}_j}, \iota) \subseteq Arc_{i+1}(\Gamma_{\mathcal{B}_j}, \sigma)$, and hence $Arc_{i+1}(\Gamma_{\mathcal{B}_j}, \sigma) = Arc_{i+1}(\Gamma_{\mathcal{B}_j}, \iota)$.

($\Rightarrow$) Suppose it follows for some $i \geq 0$. Then for any $\sigma_1 \in \Gamma(\sigma)$, as $Arc_i(\Gamma_{\mathcal{B}_j}, \sigma) = Arc_i(\Gamma_{\mathcal{B}_j}, \iota)$, we have $\mathcal{B}_{i+j}(\sigma) = \mathcal{B}_{i+j}(\iota)$, hence there exists some $\iota_1 \in \Gamma(\iota)$ such that $\iota_1 \in \mathcal{B}_{i+j+1}(\sigma_1)$. Let $\mathbb{S} = Arc_{i+1}(\Gamma_{\mathcal{B}_j}, \sigma) = Arc_{i+1}(\Gamma_{\mathcal{B}_j}, \iota)$, then $Arc_i(\Gamma_{\mathcal{B}_j}, \sigma_1) = Arc_i(\Gamma_{\mathcal{B}_j}, \iota_1) = \mathbb{S}_{\mathcal{B}_j(\sigma_1)}$. Hence $\mathcal{B}_{i+j}(\sigma_1) = \mathcal{B}_{i+j}(\iota_1)$, that is, $\Gamma_{\mathcal{B}_{i+j}}(\sigma) = \Gamma_{\mathcal{B}_{i+j}}$, hence $\mathcal{B}_{i+j+1}(\sigma) = \mathcal{B}_{i+j+1}(\iota)$.

For any $x \in X_{Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)}$, as $Arc_i(\Gamma_{\mathcal{B}_j}, \sigma^x) = Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)^x = Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)$, hence $\mathcal{B}_{i+j}(\sigma)^x = \mathcal{B}_{i+j}(\sigma^x) = \mathcal{B}_{i+j}(\sigma)$, thus we have $X_{Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)} \leq X_{\mathcal{B}_{i+j}(\sigma)}$. On the other hand, for any $x \in X_{\mathcal{B}_{i+j}(\sigma)}$, $\mathcal{B}_{i+j}(\sigma^x) = \mathcal{B}_{i+j}(\sigma)^x = \mathcal{B}_{i+j}(\sigma)$, hence $Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)^x = Arc_i(\Gamma_{\mathcal{B}_j}, \sigma^x) = Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)$, thus $x \in X_{Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)}$, and so $X_{\mathcal{B}_{i+j}(\sigma)} \leq X_{Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)}$. It follows that $X_{\mathcal{B}_{i+j}(\sigma)} = X_{Arc_i(\Gamma_{\mathcal{B}_j}, \sigma)}$. ■

Lemma 3.12 *Let $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, $(\sigma, \tau) \in Arc(\Gamma)$, $(\iota, \gamma) \in Arc(\Gamma[\mathcal{B}(\sigma), \mathcal{B}(\tau)])$ such that $\mathcal{B}_j(\sigma) = \mathcal{B}_j(\iota)$ for some integer $j \geq 1$. If $d_{j-1} = d$, then $\mathcal{B}_{j-1}(\tau) = \mathcal{B}_{j-1}(\gamma)$.*

Proof.

Since $\iota \in \mathcal{B}_j(\sigma)$, there exists some $\delta \in \Gamma(\iota) \cap \mathcal{B}_{j-1}(\tau) \neq \emptyset$. On the other hand, as $d = d_{j-1}$, $\delta \in \mathcal{B}_{j-1}(\gamma)$, hence $\mathcal{B}_{j-1}(\tau) = \mathcal{B}_{j-1}(\gamma)$.

■

Theorem 3.13 *Let $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ with $\hat{p}(\Gamma, X, \mathcal{B}) = (v, k, r, b, d, m, c, h)$, then we have:*

(1). *For any $\{\sigma, \tau\} \in E(\Gamma)$, $Com(\Gamma[\mathcal{B}(\sigma), \mathcal{B}(\tau)]) \supset Com(\Gamma[\mathcal{B}_1(\sigma), \mathcal{B}_1(\tau)]) \supset \cdots \supset Com(\Gamma[\mathcal{B}_{h-1}(\sigma), \mathcal{B}_{h-1}(\tau)])$.*

(2). *$S_i := \{Arc_i(\Gamma_{\mathcal{B}}, \sigma)\}$ is an X-symmetric orbit on $St_i^r(\Gamma_{\mathcal{B}})$, while $\Theta_i := \{(Arc_i(\Gamma_{\mathcal{B}}, \sigma), Arc_i(\Gamma_{\mathcal{B}}, \tau)) | (\sigma, \tau) \in Arc(\Gamma)\}$, is a self-paired X-symmetric orbit on $DSt_i^r(\Gamma_{\mathcal{B}})$ such that $\Gamma_{\mathcal{B}_i} \cong \Pi(\Gamma_{\mathcal{B}}, \Theta_i)$, for $i = 1, \cdots, h$.*

Proof.

(1). Let $\Psi = Com(\Gamma[\mathcal{B}(\sigma), \mathcal{B}(\tau)], \sigma)$, then for any $\iota \in V(\Psi)$ there exists some $(\sigma = \sigma_0, \tau_1, \cdots, \tau_{j-1}, \sigma_j = \iota) \in Arc_j(\Psi)$, where $j \geq 2$ is an integer. For any integer i with $1 \leq i \leq j-1$, let $\Psi_{h-1} = \Gamma[\mathcal{B}_{h-1}(\sigma_{i-1}), \mathcal{B}_{h-1}(\tau_i)]$, then as $d_{h-1} = d$, we have $\sigma_{i+1} \in \Psi(\tau_i) = \Psi_{h-1}(\tau_i)$, and hence $\sigma_{i+1} \in \mathcal{B}_{h-1}(\sigma_{i-1})$. It follows that $\iota \in \mathcal{B}_{h-1}(\sigma_j) = \mathcal{B}_{h-1}(\sigma_{j-1}) = \cdots = \mathcal{B}_{h-1}(\sigma)$.

(2). Clearly, S_1 is an X-symmetric orbit on $St_1^r(\Gamma_{\mathcal{B}})$. Suppose it follows for some $1 \leq i \leq h-1$. Let $\mathbb{S} = Arc_{i+1}(\Gamma_{\mathcal{B}}, \sigma)$. For any $(\sigma_0, \sigma_1, \cdots, \sigma_i) \in Arc_i(\Gamma, \sigma)$ such that $(\mathcal{B}(\sigma_0), \mathcal{B}(\sigma_1), \cdots, \mathcal{B}(\sigma_i)) \in \mathbb{S}(i)$, we are going to show that $|\mathbb{S}_{\mathbb{a}}| = r-1$. Clearly, $|\mathbb{S}_{\mathbb{a}}| \geq |\Gamma_{\mathcal{B}}(\sigma_i) \setminus \{\mathcal{B}(\sigma_{i-1})\}| = r-1$.

For any $(\iota_0, \iota_1, \cdots, \iota_i) \in Arc_i(\Gamma, \sigma)$ such that $\mathcal{B}(\iota_j) = \mathcal{B}(\sigma_j)$ for $j = 0, 1, \cdots, i$, by Lemma 3.12, as $\mathcal{B}_h(\sigma_0) = \mathcal{B}_h(\iota_0)$ and $d_{h-1} = d$, we have that $\mathcal{B}_{h-1}(\sigma_1) = \mathcal{B}_{h-1}(\iota_1)$, repeat this analysis we will have that $\mathcal{B}_{h-i}(\sigma_i) = \mathcal{B}_{h-i}(\iota_i)$. So $\Gamma_{\mathcal{B}_{h-i-1}}(\sigma_i)\Gamma_{\mathcal{B}_{h-i-1}}(\iota_i)$. Thus $\Gamma_{\mathcal{B}}(\sigma_i) \setminus \mathcal{B}(\sigma_{i-1}) = \Gamma_{\mathcal{B}}(\iota_i) \setminus \mathcal{B}(\iota_{i-1})$. It follows that $|\mathbb{S}_{\mathbb{a}}| = r-1$ and so S_{i+1} is an X-symmetric orbit on $St_i^r(\Gamma_{\mathcal{B}})$. Hence it follows for all $i = 1, 2, \cdots, h$.

As Γ is X-symmetric, it is not difficult to check that Θ_i is a self-paired X-symmetric orbit on $DSt_i^r(\Gamma_{\mathcal{B}})$ for $i = 1, 2, \cdots, h$.

For any $(\sigma, \tau) \in Arc(\Gamma)$, as Γ is X-symmetric, there exists some $z \in X$ such that $(\sigma, \tau)^z = (\tau, \sigma)$. Let $G_i = X_{Arc_i(\Gamma_{\mathcal{B}}, \sigma)}$, then by Lemma 3.11 $G_i = X_{\mathcal{B}_i(\sigma)}$, and hence $\Gamma_{\mathcal{B}_i} \cong Cos(X, G_i, G_i z G_i) \cong \Pi(\Gamma_{\mathcal{B}_i}, \Theta_i)$.

∎

Theorem 3.14 *Let* $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ *with* $r \geq 2$ *and* $d = 1$, *such that* Γ *is* (X, s)*-arc-transitive for some integer* $s \geq 1$. *Let* $l = \max\{s, m\}$, *then* $\Theta := \{(Arc_l(\Gamma_{\mathcal{B}}, \sigma), Arc_l(\Gamma_{\mathcal{B}}, \tau)) | (\sigma, \tau) \in Arc(\Gamma)\}$ *is a self-paired* (X, s)*-arc-transitive orbit on* $DSt_l^r(\Gamma_{\mathcal{B}})$ *such that* $X_{Arc_l(\Gamma_{\mathcal{B}}, \sigma)} \cap X_{\mathcal{B}(\tau)} = X_{Arc_l(\Gamma_{\mathcal{B}}, \tau)} \cap X_{\mathcal{B}(\sigma)}$. *Moreover,* $\Pi(\Gamma_{\mathcal{B}}, \Theta) \cong \Gamma_{\mathcal{B}_m}$ *is* (X, s)*-arc-transitive.*

Proof. As $d = 1$, we have $h = m$. Then by Theorem 3.13, $\Pi(\Gamma_{\mathcal{B}}, \Theta) \cong \Gamma_{\mathcal{B}_l} \cong \Gamma_{\mathcal{B}_m}$. Again by $d = 1$, we have $\Gamma \cong \Gamma_{\mathcal{B}_m}$ or Γ is a cover of $\Gamma_{\mathcal{B}_m}$, hence $\Gamma_{\mathcal{B}_m}$ is (X, s)-arc-transitive. ∎

4 Proof of the main results

In this section, we are going to prove Theorem 1.1.

($\Leftarrow$)

(a). If there exists a self-paired X-symmetric orbit Θ on $DSt^r(\Sigma)$, let $\Gamma = \Pi(\Sigma, \Theta)$, $S = St(\Theta)$, $\mathcal{B} = \{S_\sigma | \sigma \in V(\Sigma)\}$, where $S_\sigma := \{\mathbb{S} \in S | \mathfrak{C}_\Sigma(\mathbb{S}) = \sigma\}$ for any $\sigma \in V(\Sigma)$, then by Theorem 2.2, $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, Γ is $(X, 1)$-arc-transitive.

(b). Let $z \in X_{\{\psi,\delta\}} \setminus X_{(\sigma,\tau)} \neq \emptyset$, then $z \in X_{\{\psi,\delta\}} \setminus X_\psi$ and $z \in X_{\{\sigma,\tau\}} \setminus X_{(\sigma,\tau)}$. Thus $(\psi, \delta)^z = (\delta, \psi)$ and $(\sigma, \tau)^z = (\tau, \sigma)$. Let $G = X_\sigma$, $H = X_\psi$, $\Gamma = Cos(X, H, HzH)$, then $\Gamma \cong e\Psi$, where $e = |E(\Sigma)|$. Let $\mathcal{B} = \{[G^x : H^x] | x \in X\}$, then $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ with $r = |H : (H \cap G^z)| = 1$ and $\Gamma_{\mathcal{B}} \cong Cos(X, G, GzG) \cong \Sigma$.

(c). Let $\Gamma = \Pi(\Sigma, \Theta)$, $S = St(\Theta)$, $\mathcal{B} = \{S_\sigma | \sigma \in V(\Sigma)\}$, where $S_\sigma := \{\mathbb{S} \in S | \mathfrak{C}_\Sigma(\mathbb{S}) = \sigma\}$ for any $\sigma \in V(\Sigma)$, then by Theorem 3.6, $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$, Γ is (X, s)-arc-transitive.

($\Rightarrow$) Let $(\Gamma, X, \mathcal{B}) \in \mathcal{G}$ such that $\Gamma_\mathcal{B} \cong \Sigma$. If Γ is (X, s)-arc-transitive, then one of the following cases occurs:

(a). $s = 1$, let $\Theta = \{(\Gamma_\mathcal{B}[\sigma], \Gamma_\mathcal{B}[\tau]) | (\sigma, \tau) \in \Gamma\}$, then by Theorem 2.4, Θ is a self-paired X-symmetric orbit on $DSt^r(\Gamma_\mathcal{B})$.

(b). If $r = 1$, then by [2], $\Gamma \cong e\Gamma[\mathcal{B}(\psi), \mathcal{B}(\delta)]$ for any $\{\psi, \delta\} \in E(\Gamma)$. Let $\Psi = \Gamma[\mathcal{B}(\psi), \mathcal{B}(\delta)]$, then Ψ is bipartite, and as Γ is X-symmetric, Γ is (X, s)-arc-transitive if and only if Ψ is $(X_{V(\Psi)}, s)$-arc-transitive. Moreover, as $s \geq 2$, then $val(\Psi) \geq 2$, and so $X_\psi < X_{(\mathcal{B}(\psi), \mathcal{B}(\delta))}$, $X_{\{\psi,\delta\}} < X_{\{\mathcal{B}(\psi),\mathcal{B}(\delta)\}}$. As Γ is X-symmetric, there exists $z \in X$ such that $(\psi, \delta)^z = (\delta, \psi)$, then $(\mathcal{B}(\psi), \mathcal{B}(\delta))^z = (\mathcal{B}(\delta), \mathcal{B}(\psi))$, hence $z \in X_{\{\psi,\delta\}} \setminus X_{(\sigma,\tau)}$.

(c). As $r \geq 2$, $\mathbb{b} = b \geq r + 1 \geq 3$. Also, for any $\psi \in V(\Gamma)$, there exits $\delta, \gamma \in \Gamma(\psi)$ such that $\mathcal{B}(\delta) \neq \mathcal{B}(\gamma)$. If $d \geq 2$, then there exits $\epsilon \in \mathcal{B}(\delta) \setminus \{\delta\}$ such that $(\delta, \psi, \gamma) \neq (\delta, \psi, \epsilon) \in Arc_2(\Gamma)$, as Γ is $(X, 2)$-arc-transitive, there exists some $x \in X$ such that $(\delta, \psi, \gamma) = (\delta, \psi, \epsilon)^x$. Then $\mathcal{B}(\delta) = \mathcal{B}(\delta^x) = \mathcal{B}(\delta)^x = \mathcal{B}(\epsilon)^x = \mathcal{B}(\gamma)$, a contradiction, hence $d = 1$. Let $l = \max\{s, m+1\}$, $\Theta = \{(Arc_l(\Gamma_\mathcal{B}, \sigma), Arc_l(\Gamma_\mathcal{B}, \tau)) | (\sigma, \tau) \in Arc(\Gamma)\}$, then by Theorem 3.14, Θ is a self-paired (X, s)-arc-transitive orbit on $DSt_l^r(\Gamma_\mathcal{B})$ such that $X_{Arc_l(\Gamma_\mathcal{B},\sigma)} \cap X_{\mathcal{B}(\tau)} = X_{Arc_l(\Gamma_\mathcal{B},\tau)} \cap X_{\mathcal{B}(\sigma)}$.

References

[1] N.L.Biggs, Algebraic graph theory, Second Edition, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1993.

[2] Z.P. Lu and S. Zhou, Finite Symmetric Graphs with Two-Arc Transitive Quotients II, *J. of Graph Theory* **56**(2007), 167-193.

[3] C.H.Li, On finite $s-$transitive graphs of odd order, J.Combin.Theory Ser.B 81(2001)307-317.